

# ASPIRIN

(NSAID) • 325 mg each

PAIN RELIEVER / FEVER REDUCER

COMPARE TO THE ACTIVE  
INGREDIENT OF BAYER® ASPIRIN™



Actual Size

100 TABLETS



3 57896 90101 1

This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Bayer® Aspirin.

DISTRIBUTED BY:  
GERI-CARE PHARMACEUTICAL CORP.  
1650 63 Street, Brooklyn, NY 11204

50844  
ORG01923912  
SEAL BREAKS IF PROPERLY USED  
SEE USER'S MANUAL FOR MORE INFORMATION

## Drug Facts

### Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)\* ..... Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

### Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
- backache
- the common cold
- toothache
- premenstrual and menstrual cramps

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:
 

- facial swelling
- shock
- hives
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause a severe stomach bleeding. The chance is higher if you:
 

- are age 60 or older

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

### Drug Facts (continued)

■ have had stomach ulcers or bleeding problems  
■ have 3 or more alcoholic drinks every day while using this product. ■ take more or for a longer time than directed  
Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

**Ask a doctor before use if** ■ you have asthma  
■ stomach bleeding warning applies to you  
■ you have a history of stomach problems, such as heartburn  
■ you have high blood pressure, heart disease, liver cirrhosis or kidney disease ■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are taking a prescription drug for** ■ gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint  
■ vomit blood ■ have bloody or black stools  
■ have stomach pain that does not get better  
■ an allergic reaction occurs. Seek medical help right away.  
■ ringing in the ears or loss of hearing occurs  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present ■ new symptoms occur  
These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.**

**Directions** ■ drink a full glass of water with each dose  
■ adults and children 12 years and over: take 1 to 2 tablets every 4 hours while symptoms last. Do not take more than 12 tablets in 24 hours unless directed by a doctor.  
■ children under 12 years: do not use unless directed by a doctor

### Other information

- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

**Inactive ingredient** cinn starch

**Questions or comments?** 1-800-540-3765

STOP PEELING