

Drug Facts

Active ingredient (in each caplet)

Loperamide HCl 2 mg.....Anti-diarrheal

Purpose

Uses • controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Do not use • if you have bloody or black stool

Ask a doctor before use if you have • fever • mucus present in your stool • a history of liver disease

Ask a doctor or pharmacist before use if you are • taking antibiotics

When using this product • tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if • diarrhea lasts for more than 2 days • symptoms get worse • you get abdominal swelling. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

NDC 57896-381-48



Anti-Diarrheal

Loperamide HCl

Controls the symptoms of Diarrhea

48 Caplets
2 mg each

*Compare to the active ingredient in IMODIUM A-D®

Tamper Evident: Do not use if imprinted seal under cap is broken or missing



Directions: drink plenty of clear fluids to help prevent dehydration, which may accompany diarrhea • do not exceed recommended dose

Adults and children 12 years and over

2 caplets after the first loose bowel movement; 1 caplet after each subsequent loose bowel movement; do not exceed 4 caplets in 24 hours

Children under 12 years

Ask a doctor

Other information

• store at controlled room temperature, 20°-25°C (68°-77°F)

Inactive ingredients: cellulose, corn starch, D&C Yellow #10, dicalcium phosphate, FD&C Blue #1, magnesium stearate, silica gel

Questions or comments? call 1-800-540-3765

*This product is not manufactured or distributed by the owner of the registered trademark IMODIUM®. REV 0613

Dist. by: Geri-Care
Pharmaceuticals Corp.
1650 63rd Street
Brooklyn, NY 11204 3

