

3 57896 22601 5



100 Caplets

Compare to active ingredients in **EXCEDRIN**®\*

**EXTRA STRENGTH**  
**PAIN-RELIEVER**  
 Acetaminophen 250 mg  
 Aspirin 250 mg (NSAID)  
 Caffeine 65 mg

GERI CARE®  
 NDC 57896-216-01

### Drug Facts

**Tamper Evident:** Do not use if imprinted seal under cap is missing or broken.

### Active ingredients (in each caplet)

Acetaminophen 250 mg.....	Pain reliever
Aspirin 250 mg (NSAID-nonsteroidal anti-inflammatory drug).....	Pain reliever
Caffeine 65 mg.....	Pain reliever aid

### Purpose

**Uses** • temporarily relieves minor aches and pains

### Warnings

**Reye's Syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives • facial swelling • asthma (wheezing) • shock

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • rash • skin reddening • blisters If a skin reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • are age 60 or older • have 3 or more alcoholic drinks every day while using this product

• take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

**Do not use** • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if** • you have liver disease • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma

**Ask a doctor or pharmacist before use if you are** • taking a prescription drug for diabetes, gout, or arthritis • taking any other drug or are under a doctor's care for any serious condition

**Stop use and ask a doctor if** • an allergic reaction occurs. Seek medical help right away.

- you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs • redness or swelling is present
- pain gets worse or lasts for more than 10 days • fever gets worse or lasts for more than 3 days • new symptoms occur These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions • do not take more than directed

- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours. Do not take more than 8 caplets in 24 hours. • children under 12: ask a doctor

### Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

**Inactive ingredients** corn starch, crospovidone, hypromellose, microcrystalline cellulose, povidone, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

### Questions or comments? 1-800-540-3765

Dist. By: Gerl-Care Pharmaceuticals Corp. 1650 63rd St., Brooklyn, NY 11204

\*This product is not manufactured or distributed by the owner of the registered trademark Excedrin®.

REV  
0220L