

Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*.....Pain reliever
*nonsteroidal anti-inflammatory drug

Purpose

Uses • for the temporary relief of minor aches and pains • ask your doctor about other uses for aspirin

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if • the stomach bleeding warning applies to you • you are taking a diuretic • you have a history of stomach problems, such as heartburn • you have: ◦ high blood pressure ◦ heart disease ◦ liver cirrhosis ◦ kidney disease ◦ asthma

Ask a doctor or pharmacist before use if you are • taking any other drug containing an NSAID (prescription or nonprescription) • taking a blood thinning (anticoagulant) or steroid drug • taking a prescription drug for diabetes, gout or arthritis ▶

GERI-CARE®

NDC 57896-921-10

Regular Strength Enteric Coated Aspirin

Pain Reliever (NSAID)

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

1000 Tablets 325 mg each

Stop use and ask a doctor if • you experience any of the following signs of stomach bleeding: ◦ feel faint ◦ vomit blood ◦ have bloody or black stools ◦ have stomach pain that does not get better • an allergic reaction occurs. Seek medical help right away. • pain gets worse or lasts more than 10 days • redness or swelling is present • fever gets worse or lasts more than 3 days • any new symptoms occur • ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • drink a full glass of water with each dose • swallow whole, do not chew or crush • do not exceed recommended dose

adults and children 12 years and older	take 1-2 tablets every 4 hours, as needed, not more than 12 tablets in 24 hours, or as directed by a doctor
children under 12	ask a doctor

Other information • store at 20°C - 25°C (68°F - 77°F);
excursions permitted between 15°C - 30°C (59°F - 86°F)

Inactive ingredients: black iron oxide, cellulose, corn starch, D&C yellow #10 lake, FD&C yellow #6 lake, hypromellose, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, propylene glycol, shellac wax, silicon dioxide, simethicone, sodium alginate, sodium bicarbonate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate.

Questions or comments? 1-800-540-3765

DIST. BY: GERI-CARE PHARMACEUTICALS CORP.
1650 63 Street, Brooklyn, New York 11204

REV 0521L

