



NDC 57896-951-05

THE SENSIBLE CHOICE FOR PRICE AND HEALTH.

NAPROXEN SODIUM

Pain Reliever / Fever Reducer

(NSAID) nonsteroidal anti-inflammatory drug



Compare to the active ingredient in ALEVE®*

50 Tablets
220mg Each

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active ingredient (in each tablet)

Naproxen sodium 220 mg
(Naproxen 200 mg) (NSAID)* Pain Reliever/
*nonsteroidal anti-inflammatory drug Fever Reducer

Purposes

- Uses** ■ Temporarily relieves minor aches and pains due to:
- backache
 - headache
 - menstrual cramps
 - minor pain of arthritis
 - muscular aches
 - toothache
 - the common cold
- Temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- asthma (wheezing)
- hives
- rash
- shock
- skin reddening
- blisters
- facial swelling
- skin reddening

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

*This product is not manufactured or distributed by the owner of the registered trademark Aleve®.

DIST. BY: GERI-CARE PHARMACEUTICALS CORP.

1650 63rd Street, Brooklyn, NY 11204 Made in India



3

57896 95105

4

PEEL HERE FOR MORE DRUG FACTS

REV 0316

Questions or comments?
Call 1-800-540-3765

STOP PEELING

Inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, FD&C blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, titanium dioxide.

Other information

- each tablet contains: sodium 20 mg ■ store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F) ■ If side effects occur, you may report side effects to FDA at 1-800-FDA-1088

■ children under 12 years: ask a doctor

do not exceed 2 tablets in any 8 to 12 hour period

do not exceed 3 tablets in a 24 hour period

for the first dose you may take 2 tablets within the first hour

■ adults and children 12 years and older: take 1 tablet every 8 to 12 hours while symptoms last

■ the smallest effective dose should be used

■ drink a full glass of water with each dose

Directions ■ do not take more than directed

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

- any new symptoms appear

■ you have difficulty swallowing

■ it feels like the pill is stuck in your throat

■ redness or swelling is present in the painful area

■ pain gets worse or lasts more than 10 days

■ fever gets worse or lasts more than 3 days

■ you have difficulty swallowing

■ you have difficulty swallowing

■ it feels like the pill is stuck in your throat

■ redness or swelling is present in the painful area

■ any new symptoms appear

■ you experience any of the following signs of stomach bleeding: feel faint vomit blood

have bloody or black stools

have stomach pain that does not get better

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

you have difficulty swallowing

you have difficulty swallowing

it feels like the pill is stuck in your throat

redness or swelling is present in the painful area

any new symptoms appear

Drug Facts (continued)

When using this product

■ take with food or milk if stomach upset occurs

■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding: feel faint vomit blood

have bloody or black stools

have stomach pain that does not get better

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

you have difficulty swallowing

you have difficulty swallowing

it feels like the pill is stuck in your throat

redness or swelling is present in the painful area

any new symptoms appear