

REGULAR STRENGTH
ACETAMINOPHEN
325 mg
 PAIN RELIEVER/ FEVER REDUCER



Compare to the active ingredient
 in Regular Strength **TYLENOL®***

50 Tablets

TAMPER EVIDENT: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active ingredient (in each tablet) **Purposes**
 Acetaminophen 325 mg Pain Reliever/
 Fever Reducer

- Uses**
- temporarily relieves minor aches and pains
 - temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg (12 tablets) of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

- Do not use**
- with any other drug containing acetaminophen (prescription or nonprescription).
 - if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 - if you are allergic to acetaminophen or any of the inactive ingredients in this product →

* This product is not manufactured or distributed by the owner of the registered trademark **TYLENOL®**.

DIST. BY: **GERI-CARE PHARMACEUTICALS CORP.**
 1295 Towbin Ave, Lakewood, NJ 08701

REV 0922AA

 3 57896 10105 3

PEEL HERE FOR MORE DRUG FACTS

Drug Facts (continued)

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- adults and children 12 years and over: take 2 tablets every 4-6 hours, as needed; not more than 10 tablets in 24 hours. Do not take for more than 10 days unless directed by a doctor.
- children under 12 years: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)

Inactive ingredients: povidone, sodium starch glycolate, starch, stearic acid.

Questions or comments? 1-800-540-3765

STOP PEELING

ADHESIVE