

## Drug Facts

### Active ingredient (in each tablet)      Purposes

Acetaminophen 500 mg .....Pain Reliever /  
Fever Reducer

**Uses** • temporarily relieves minor aches and pains  
• temporarily reduces fever

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 tablets (4,000 mg of acetaminophen) in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away.

**Do not use** • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
• if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have** liver disease.

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin.

**Stop use and ask a doctor if** • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present  
These could be signs of a serious condition. ▶



EXTRA STRENGTH

**ACETAMINOPHEN**  
**500 mg**

**PAIN RELIEVER/ FEVER REDUCER**



Compare to the active ingredient in  
Extra Strength **TYLENOL®** Tablets\*

**1000 Tablets**

NDC 57896-201-10

## Drug Facts (continued)

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions • do not take more than directed

Adults and children 12 years of age and over	Take 2 tablets every 6 hours, as needed; not more than 6 tablets in 24 hours. Do not take for more than 10 days unless directed by a doctor.
Children under 12 years	ask a doctor

**Other information • TAMPER EVIDENT:** Do not use if imprinted seal under cap is missing or broken.  
• store at 20°C-25°C (68°F-77°F)

**Inactive ingredients:** povidone, sodium starch glycolate, starch, stearic acid.

### Questions or comments? 1-800-540-3765

\*This product is not manufactured or distributed by the owner of the registered trademark **TYLENOL®**.

DIST. BY: **GERI-CARE** PHARMACEUTICALS CORP.

1295 Towbin Ave  
Lakewood, NJ 08701

REV 1123A

