

Drug Facts

Active ingredient (in each tablet) Purposes

Acetaminophen 500 mg Pain Reliever /
Fever Reducer

Uses • temporarily relieves minor aches and pains
• temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 8 tablets (4,000 mg of acetaminophen) in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening
• blisters • rash If a skin reaction occurs, stop use and seek medical help right away.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
• if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • new symptoms occur
• redness or swelling is present
These could be signs of a serious condition. ►

NDC 57896-219-10



EXTRA STRENGTH ACETAMINOPHEN 500 mg PAIN RELIEVER/ FEVER REDUCER



Compare to the active ingredient in
Extra Strength **TYLENOL®** Tablets*

1000 Tablets

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • do not take more than directed

Adults and children
12 years of age
and over

Take 2 tablets every 6 hours, as
needed; not more than 6 tablets
in 24 hours. Do not take for
more than 10 days unless
directed by a doctor.

Children under 12 years ask a doctor

Other information • TAMPER EVIDENT: Do not use if imprinted seal under cap is missing or broken.
• store at 20°C-25°C (68°F-77°F)

Inactive ingredients: povidone, sodium starch
glycolate, starch, stearic acid.

Questions or comments? 1-800-540-3765

*This product is not manufactured or distributed by the owner of the registered trademark **TYLENOL®**.

DIST. BY: **GERI-CARE PHARMACEUTICALS CORP.**

1295 Towbin Avenue
Lakewood, NJ 08701

REV 1124AAIG

3 57896 21910 9