



EXTRA STRENGTH NIGHT TIME

PAIN MEDICINE

Acetaminophen 500 mg

Diphenhydramine HCl 25 mg

PAIN RELIEVER & SLEEP AID



Compare to the active
ingredients in **TYLENOL® PM***

50 Caplets

TAMPER EVIDENT: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active ingredients (in each caplet) Purposes
Acetaminophen 500 mg Pain Reliever
Diphenhydramine HCl 25 mg Sleep-aid

Uses • for the temporary relief of occasional headaches and minor aches and pains along with accompanying sleeplessness

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take
 • more than 4,000 mg of acetaminophen in 24 hours
 • with other drugs containing acetaminophen
 • 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
 • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 • with any other product containing diphenhydramine, even one used on skin in children under 12 years of age
 • if you have ever had an allergic reaction to this product or any of its ingredients →

* This product is not manufactured or distributed by the owner of the registered trademark **TYLENOL® PM**.

Dist. By: **Gert-Care Pharmaceuticals Corp.**
 1295 Towbin Ave, Lakewood, NJ 08701

REV 08/24/01



3 57896 22405 9

Keep out of reach of children

Drug Facts (continued)

Ask a doctor before use if you have
 • liver disease • glaucoma • a breathing problem such as emphysema or chronic bronchitis • trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking • the blood thinning drug warfarin • sedatives or tranquilizers

When using this product • drowsiness will occur • avoid alcoholic drinks • do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if
 • sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness. • pain gets worse or lasts for more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur

These may be signs of a serious condition. **If pregnant or breast-feeding**, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • do not take more than directed (see overdose warning)

• adults and children 12 years and over: take 2 caplets at bedtime. Do not take more than 2 caplets of this product in 24 hours.
 • children under 12 years: do not use

Other information • store at controlled room temperature 20-25 °C (68-77° F)

Inactive ingredients: croscarmellose sodium, FD&C blue #1 lake, hypromellose, microcrystalline cellulose, PEG, povidone, silica, starch, stearic acid, titanium dioxide.

Questions or comments? 1-800-540-3765

STOP PEELING