



**EXTRA STRENGTH  
NIGHT TIME  
PAIN MEDICINE**  
Acetaminophen 500 mg  
Diphenhydramine HCl 25 mg  
PAIN RELIEVER & SLEEP AID



Compare to the active  
ingredients in **TYLENOL® PM\***

**100 Caplets**

**TAMPER EVIDENT: Do not use if imprinted seal under cap is missing or broken.**

### Drug Facts

**Active ingredients (in each caplet)** *Purposes*  
Acetaminophen 500 mg.....Pain Reliever  
Diphenhydramine HCl 25 mg.....Sleep-aid

**Uses** • for the temporary relief of occasional headaches and minor aches and pains along with accompanying sleeplessness

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash if a skin reaction occurs, stop use and seek medical help right away.

**Do not use** • with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains

- acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- in children under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients →

\* This product is not manufactured or distributed by the owner of the registered trademark **TYLENOL® PM**.

Dist. By: **Geri-Care Pharmaceuticals Corp.**  
1295 Towbin Ave, Lakewood, NJ 08701  
Product of India  
REV 06/22B



3 57896 22401 1

**PEEL HERE FOR MORE DRUG FACTS** →

### Drug Facts (continued)

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are taking** • the blood thinning drug warfarin • sedatives or tranquilizers

**When using this product** • drowsiness will occur • avoid alcoholic drinks • do not drive a motor vehicle or operate machinery

**Stop use and ask a doctor if**

- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These may be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

### Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** • do not take more than directed (see overdose warning)

- adults and children 12 years and over: take 2 caplets at bedtime. Do not take more than 2 caplets of this product in 24 hours.
- children under 12 years: do not use

**Other information** • store at controlled room temperature 20-25 °C (68-77° F)

**Inactive ingredients:** cellulose, croscarmellose sodium, FD&C blue #1 lake, FD&C blue #2 lake, hypromellose, magnesium stearate, PEG, polyvinyl alcohol, polydione, purified water, silicon dioxide, sodium starch glycolate, starch, talc, titanium dioxide

**Questions or comments? 1-800-540-3765**

**STOP PEELING**