

265.500

LOT/EXP

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING



**Drug Facts**

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

**Active ingredient (in each caplet)**

Loperamide HCl 2 mg

**Purpose**

Anti-diarrheal

**Use** controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.

**Drug Facts (continued)**

**Heart alert:** Taking more than directed can cause serious heart problems or death.

**Do not use** if you have bloody or black stool.

**Ask a doctor before use if you have** ■ a fever ■ mucus in the stool ■ a history of liver disease ■ a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product,** tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if** ■ symptoms get worse ■ diarrhea lasts for more than 2 days ■ you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea  
■ find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

**Other information**

■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN  
■ store between 20°-25°C (68°-77°F) ■ see end flap for expiration date and lot number

**Inactive ingredients** corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

**Questions or comments?** Call 1-800-540-3765

1. To open, tear at perforations and remove a section.  
2. Use to cut through plastic for caplet.



\*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Imodium® A-D.  
50844 REV0619A37508  
REV 0321

DISTRIBUTED BY:  
**GERI-CARE PHARMACEUTICAL CORP.**  
1650 63 Street, Brooklyn, NY 11204

B-0617-375-08  
REV0619A37508



**ANTI-DIARRHEAL**

NDC 57896-381-25



**ANTI-DIARRHEAL**  
Loperamide HCl Tablets, 2 mg  
Anti-Diarrheal

Controls the  
Symptoms of Diarrhea



actual size

compare to active  
ingredient in Imodium® A-D\*

24 caplets

LEBG81 6D C7359