

### Drug Facts (continued)

- Directions** • take with a glass of water  
• **do not take more than directed**  
• adults and children 12 years and older:  
take 1 to 3 tablets in a single daily dose  
• children 12 and under: ask a doctor

### Other information

- store at controlled room temperature 20°-25°C (68°-77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid excessive humidity
- package not child resistant

**Inactive ingredients:** acacia, ammonium hydroxide, calcium carbonate, corn starch, D&C yellow #10 lake, FD&C yellow #6 lake, hydroxyethylcellulose, iron oxide black, lactose, magnesium stearate, methylparaben, PEG, polydextrose, polyvinyl acetate phthalate, propylparaben, propylene glycol, povidone, shellac, simethicone, silica, sodium alginate, sodium benzoate, sodium bicarbonate, stearic acid, sucrose, talc, titanium dioxide, triacetin, triethyl citrate, wax.

### Questions or comments? 1-800-540-3765

STOP PEELING

VALUE SIZE

NDC 57896-441-20

GERI CARE®



# BISACODYL

## STIMULANT LAXATIVE

### Enteric Coated Tablets

Compare to active ingredient in **Dulcolax® Laxative\***

200 Tablets 5 mg each

**Tamper Evident:** Do not use if imprinted seal under cap is missing or broken.

## Drug Facts

**Active ingredient (in each tablet)** Purpose  
Bisacodyl 5 mg.....Stimulant Laxative

**Uses** • for relief of occasional constipation and irregularity • this product generally produces bowel movement in 6 to 12 hours

### Warnings

**Do not use** if you cannot swallow without chewing.

**Ask a doctor before use if you have**

- stomach pain, nausea, vomiting
- noticed a sudden change in bowel habits that lasts more than 2 weeks

**When using this product** • do not chew or crush tablet(s) • do not use within 1 hour after taking an antacid or milk • it may cause stomach discomfort, faintness and cramps

**Stop use and ask a doctor if** • you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition. • you need to use a laxative for more than 1 week

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

\*This product is not manufactured or distributed by the owner registered trademark DULLCOLAX®.

Dist. by: **Geri-Care Pharmaceuticals Corp.**  
1295 Towbin Avenue, Lakewood, NJ 08701



REV 1023L

PEEL HERE FOR MORE DRUG FACTS →