

# ASPIRIN

## 325 mg (NSAID)

### PAIN RELIEVER / FEVER REDUCER

### REGULAR STRENGTH

Compare to active ingredient  
in **BAYER® Aspirin™**

Actual Size

100 Tablets



3 5789690101

...This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Aspirin.

REVO122024912  
DISTRIBUTED BY: GERI-CARE PHARMACEUTICAL CORP.

1295 Towbin Ave., Lakewood, NJ 08701

## Drug Facts

### Active Ingredient (in each tablet)

Aspirin 325 mg (NSAID) ..... Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses** temporarily relieves:

- pain and fever of colds
- headache
- muscle pain
- toothache
- menstrual pain

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if chills or in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- shock
- hives
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ have 3 or more alcoholic drinks every day while using this product  
■ take more or for a longer time than directed

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED STRIP SEAL UNDER CAP IS BROKEN OR MISSING**

PEEL HERE FOR MORE DRUG FACTS

Adhesive Only

STOP PEELING

Questions or comments? 1-800-340-3765

**Inactive ingredient** corn starch

**Other information** ■ use by expiration date on package  
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

**Directions** ■ **do not take more than directed**  
■ drink a full glass of water with each dose

■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor

■ children under 12 years: do not use unless directed by a doctor

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**These could be signs of a serious condition.**

■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present  
■ new symptoms occur

Seek medical help right away.

**Stop use and ask a doctor if** ■ an allergic reaction occurs.

■ you experience any of the following signs of stomach bleeding: vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better

■ ringing in the ears or a loss of hearing occurs  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present  
■ new symptoms occur

**Ask a doctor or pharmacist before use if you are taking a prescription drug for** ■ gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if** ■ an allergic reaction occurs.

■ you experience any of the following signs of stomach bleeding: vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better

■ ringing in the ears or a loss of hearing occurs  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present  
■ new symptoms occur

**Ask a doctor or pharmacist before use if you are taking a prescription drug for** ■ gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if** ■ an allergic reaction occurs.

■ you experience any of the following signs of stomach bleeding: vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better

## Drug Facts (continued)

■ have had stomach ulcers or bleeding problems  
■ have a blood thinning (anticoagulant) or steroid drug  
■ do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer  
■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if** ■ you have asthma  
■ you have a history of stomach problems, such as heartburn  
■ you have high blood pressure, heart disease, liver condition, or kidney disease  
■ you are taking a diuretic