

ASPIRIN

325 mg (NSAID)

PAIN RELIEVER / FEVER REDUCER
REGULAR STRENGTH



Actual Size

Compare to active ingredient
in **BAYER®** Aspirin**

1000 Tablets

Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)* Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Purpose

Uses

temporarily relieves: ■ pain and fever of colds ■ headache
■ toothache ■ muscle pain ■ minor pain of arthritis ■ menstrual pain

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older

■ take a blood thinning (anticoagulant) or steroid drug

■ have 3 or more alcoholic drinks every day while using this product

■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ take more or for a longer time than directed

■ have had stomach ulcers or bleeding problems

Do not use

■ if you are allergic to aspirin or any other pain reliever/fever reducer

■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

■ you have asthma ■ you are taking a diuretic

■ stomach bleeding warning applies to you

■ you have a history of stomach problems, such as heartburn

■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ➔

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug for

■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if

■ an allergic reaction occurs. Seek medical help right away.

■ you experience any of the following signs of stomach bleeding:

■ have bloody or black stools ■ feel faint

■ have stomach pain that does not get better ■ vomit blood

■ ringing in the ears or a loss of hearing occurs

■ pain gets worse or lasts more than 10 days ■ redness or swelling is present

■ fever gets worse or lasts more than 3 days ■ new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ do not take more than directed

■ drink a full glass of water with each dose

■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor

■ children under 12 years: do not use unless directed by a doctor

Other information

■ use by expiration date on package

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredient

corn starch

Questions or comments? 1-800-540-3765


50844 REV0122D24916

**This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Aspirin.

DISTRIBUTED BY:

GERI-CARE PHARMACEUTICAL CORP.

1295 Towbin Ave., Lakewood, NJ 08701 3



57896 90110 3