

NDC 57896-901-10



ASPIRIN

(NSAID) • 325 mg each

PAIN RELIEVER / FEVER REDUCER

COMPARE TO THE ACTIVE INGREDIENT OF BAYER® ASPIRIN™

1 000 TABLETS



Actual Size

Drug Facts

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purpose

Uses

temporarily relieves minor aches and pains due to: ■ muscular aches
■ headache ■ minor pain of arthritis ■ toothache ■ backache
■ premenstrual and menstrual cramps ■ the common cold

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you have asthma ■ you are taking a diuretic

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding: ■ vomit blood ■ feel faint ■ have bloody or black stools
 - have stomach pain that does not get better
 - an allergic reaction occurs. Seek medical help right away.
 - ringing in the ears or loss of hearing occurs ■ redness or swelling is present
 - pain gets worse or lasts more than 10 days ■ new symptoms occur
 - fever gets worse or lasts more than 3 days
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 to 2 tablets every 4 hours while symptoms last. Do not take more than 12 tablets in 24 hours unless directed by a doctor.
- children under 12 years: do not use unless directed by a doctor

Other information

- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredient

corn starch

Questions or comments? 1-800-540-3765

This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Bayer® Aspirin. 50844 ORG011324916

DISTRIBUTED BY:
GERI-CARE PHARMACEUTICAL CORP.
1650 63 Street, Brooklyn, NY 11204

