

GERI CARE®

ADULT LOW DOSE
CHEWABLE**ASPIRIN 81 mg**
PAIN RELIEVER (NSAID)Compare to active ingredient in
AYER® Low Dose Aspirin*

36 Tablets

Drug Facts

Active ingredient (in each tablet)
Aspirin 81 mg (NSAID)
*nonsteroidal anti-inflammatory drug

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

Purpose
Pain reliever

Uses for the temporary relief of minor aches and pains or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ snuck ■ asthma (wheezing).

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ this product for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

* This product is not manufactured or distributed by the owner of the registered trademark Bayer®. Dist. by: **Geri-Care Pharmaceuticals Corp.**
1295 Townin Avenue, Lakewood, NJ 08701

REV 1123



3 57896 91136 2
PEEL HERE FOR MORE FACTS →

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug for
■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stool
■ have stomach pain that does not get better
■ pain gets worse or lasts more than 10 days
■ redness or swelling is present
■ new symptoms occur
■ ringing in the ears or loss of hearing occurs

if pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

■ drink a full glass of water with each dose
■ do not exceed recommended dose

adults and children take 4-8 tablets every 4 hours, as needed, not more than 48 tablets in 24 hours, or as directed by a doctor

children under 12 ask a doctor

Other information ■ store at room temperature 15°C - 30°C (59°F - 86°F)

Inactive ingredients: corn starch, dextrates, FD&C yellow #6 aluminum lake, orange flavor, saccharin sodium.

Questions or comments? 1-800-540-3765

STOP PEELING