



REGULAR STRENGTH
ENTERIC COATED

ASPIRIN
325 mg

PAIN RELIEVER (NSAID)

100 Tablets

TAMPER EVIDENT: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active ingredient (in each tablet) *Purpose*
Aspirin, 325 mg (NSAID)Pain reliever
nonsteroidal anti-inflammatory drug

Uses ■ for the temporary relief of minor aches and pains
■ ask your doctor about other uses for aspirin

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ the stomach bleeding warning applies to you ■ you are taking a diuretic ■ you have a history of stomach problems, such as heartburn ■ you have: ■ high blood pressure ■ heart disease ■ liver cirrhosis ■ kidney disease ■ asthma

Dist. by: GERI-CARE PHARMACEUTICALS CORP.
1295 Towbin Ave, Lakewood, NJ 08701

REV 0722L



PEEL HERE FOR MORE DRUG FACTS →

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, gout or arthritis.

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away, ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ pain gets worse or lasts more than 10 days ■ stomach pain that does not get better ■ redness or swelling is present ■ fever gets worse or lasts more than 3 days ■ any new symptoms occur ■ ringing in the ears or loss of hearing occurs These can be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not exceed recommended dose ■ swallow whole; do not chew or crush ■ drink a full glass of water with each dose

adults and children 12 years and older	take 1 to 2 tablets every 4 hours, as needed, not more than 12 tablets in 24 hours, or as directed by a doctor
children under 12	do not use unless directed by a doctor

Other information

■ store at 20-25°C (68-77°F); excursions permitted between 15°C - 30°C (59°F - 86°F)

Inactive ingredients: black iron oxide, cellulose, corn starch, D&C yellow #10 lake, FD&C yellow #6 lake, hydroxypropylcellulose, polyethylene glycol, polyvinyl acetate phthalate, polyethylene glycol, shellac wax, silica, stearic acid, sodium alginate, sodium bicarbonate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate.

Questions or comments? 1-800-540-3765

STOP PEELING

Spot UV