

IBUPROFEN 200 mg

PAIN RELIEVER/ FEVER REDUCER
(NSAID)



Compare to the active
ingredient in **ADVICE**®*

100 Tablets

Warning: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active ingredient (in each brown tablet) Purpose

Ibuprofen USP 200 mg (NSAID)* Pain reliever/fever reducer
Nonsteroidal anti-inflammatory drug

Uses ■ Temporarily relieves minor aches and pains due to
 □ headache □ menstrual cramps □ joint aches
 □ backache □ muscle aches □ rheumatoid
 □ minor pain of arthritis ■ Temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 ■ shock ■ facial swelling ■ asthma (wheezing) ■ skin redness
 ■ hives ■ rash ■ blisters If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

■ take more or for a longer time than directed
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ are age 60 or older ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 ■ have had stomach ulcers or bleeding problems
 ■ have 3 or more alcoholic drinks every day while using this product
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ right before or after heart surgery
 ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers

*This product is not manufactured or distributed by the owner of the registered trademark **ADVICE**®.
 Distributed by: **GERI-CARE PHARMACEUTICALS CORP.**
 1295 Towbin Ave, Lakewood, NJ 08701

REV 11/22



PEEL HERE FOR MORE DRUG FACTS

Spot UV

Drug Facts (continued)

■ you have a history of stomach problems, such as heartburn
 ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are

■ under a doctor's care for any serious condition
 ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

When using this product

■ take with food or milk, if stomach upset occurs

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: □ feel faint □ have bloody or black stools □ vomit blood □ have stomach pain that does not get better

■ you have symptoms of heart problems or stroke:
 □ chest pain □ trouble breathing □ leg swelling □ slurred speech □ weakness in one part or side of body
 ■ pain gets worse or lasts for more than 10 days
 ■ fever gets worse or lasts for more than 3 days
 ■ redness or swelling is present in the painful area
 ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ do not take more than directed
 ■ the smallest effective dose should be used
 ■ adults and children 12 years and over:

■ take 1 tablet every 4 to 6 hours while symptoms persist
 ■ pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not take more than 6 tablets in 24 hours unless directed by a doctor

■ children under 12 years: ask a doctor

Other information ■ store between 20°-25°C (68°-77°F) ■ avoid excessive heat 40°C (104°F)

Inactive Ingredients: Carnauba wax, cellulose, colloidal silicon dioxide, corn starch, hydroxypropylcellulose, lactose, magnesium stearate, polydextrose, PEG, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide.

STOP PEELING

Questions or comments? Call 1-800-540-3765