



THE SENSIBLE CHOICE
FOR PRICE AND HEALTH

NDC 57896-941-20

IBUPROFEN

PAIN RELIEVER/FEVER REDUCER (NSAID)



COMPARE TO THE ACTIVE
INGREDIENT IN ADVIL

200 Tablets
200 mg Each

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active Ingredient (in each brown tablet) *Purpose*

Ibuprofen USP, 200 mg (NSAID)* Pain reliever/
nonsteroidal anti-inflammatory drug Fever reducer

Uses

- Temporarily relieves minor aches and pains due to:
 - muscular aches
 - headache
 - backache
 - menstrual cramps
 - the common cold
 - minor pain of arthritis
- Temporarily reduces fever

Warnings

Allergists alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- shock
- facial swelling
- asthma/wheezing
- hives
- rash
- blisters
- skin redness/itching

 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing aspirin (or other nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) product)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use if

- you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers

*This product is not manufactured or distributed by the owner of the registered trademark Advil®.

Distributed by: **Geri-Care Pharmaceuticals Corp.**
1650 63rd Street, Brooklyn, NY 11204

REV 09/21



3 57896 94120 8

PEEL HERE FOR MORE DRUG FACTS

Drug Facts (continued)

■ You have a history of stomach problems, such as heartburn

■ You have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

■ You are taking a diuretic

Ask a doctor or pharmacist before use if you are

■ under a doctor's care for any serious condition

■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin

■ taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding: a foul faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better

■ You have symptoms of heart problems or stroke

■ Chest pain □ trouble breathing □ leg swelling

■ Sudden speech □ weakness in one part or side of body

■ pain gets worse or lasts for more than 10 days

■ fever gets worse or lasts for more than 3 days

■ redness or swelling is present in the painful area

■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than directed

the smallest effective dose should be used

■ adults and children 12 years and over:

□ take 1 tablet every 4 to 6 hours while symptoms persist

□ pain or fever does not respond to 1 tablet, 2 tablets may be used, □ do not take more than 6 tablets in 24 hours unless directed by a doctor

■ children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)
- avoid excessive heat, 40°C (104°F)

Inactive ingredients: carmelum wax, colloidal silicon dioxide, cellulose, corn starch, hypromellose, lactose, magnesium stearate, polydextrose, PEGs, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide.

Questions or comments? Call 1-800-540-3765

STOP PEELING