

NAPROXEN

NAPROXEN SODIUM 220 MG
PAIN RELIEVER/
FEVER REDUCER (NSAID)
Strength to last 12 hours

Compare to active ingredient
in **ALEVE®** Tablets*

100 Film Coated Tablets

Tempor Evident: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active ingredient (in each tablet)	Purpose
Naproxen sodium 220 mg	Pain Reliever
Naproxen 200 mg (NSAID)*	Fever Reducer
Nonsteroidal anti-inflammatory drug	

Uses ■ temporarily relieves minor aches and pains due to:

- headache
- toothache
- backache
- muscular aches
- menstrual cramps
- the common cold
- minor pain of arthritis

■ temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- asthma (wheezing)
- skin reddening
- shock
- blisters
- rash
- facial swelling
- hives

If an allergic reaction occurs, stop use and seek medical help right away. May cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant or steroid) drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have a more acidic stomach every day while using this product
- take more or for a longer time than directed

Heart Attack and Stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

*This product is not manufactured or distributed by the owner of the registered trademark Aleve®.
DST: BY: GERI-CARE PHARMACEUTICALS CORP.
1295 Towbin Ave. Lakewood, NJ 08701



REV 0422L

3 57896 95101 6
PEEL HERE FOR MORE DRUG FACTS

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking any other drug

When using this product

- take with food or milk, if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke
 - leg swelling
 - chest pain
 - slurred speech
 - trouble breathing
- weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing ■ it feels like the pill is stuck in your throat ■ any new symptoms appear
- redness or swelling is present in the painful area

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ do not take more than directed

- the smallest effective dose should be used
- drink a full glass of water with each dose
- adults and children 12 years and older
 - take 1 tablet every 8 to 12 hours while symptoms last
 - for the first dose you may take 2 tablets within the first hour
- do not exceed 2 tablets in any 8 to 12 hour period
- do not exceed 3 tablets in a 24 hour period
- children under 12 years: ask a doctor

Other information

- each tablet contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients

croscarmellose sodium, FdC blue #2 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, talc, titanium dioxide

Questions or comments? Call 1-800-540-5785

STOP PEELING