

NAPROXEN

NAPROXEN SODIUM 220 MG
PAIN RELIEVER/
FEVER REDUCER (NSAID)
Strength to last 12 hours

Compare to active ingredient
in **ALEVE®** Tablets*

100 Film Coated Tablets

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking any other drug ■ taking aspirin for heart attack or stroke, because naproxen may decrease the benefit of aspirin

When using this product

Mix with food or milk if stomach upset occurs

Stop use and ask a doctor if ■ you have the following signs of stomach

bleeding: ■ feel faint ■ vomit blood

■ have bloody or black stools

■ have stomach pain that does not get better

■ you have symptoms of heart problems or stroke

■ leg swelling

■ dizziness

■ weakness in one part of your body

■ painless weight loss more than 10 days

■ fever lasts more than 3 days

■ you have difficulty swallowing

■ it feels like the pill is stuck in your throat

■ any new symptoms appear

■ redness or swelling is present in the painful area

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen

sodium at 20 weeks or later in pregnancy unless definitely

ordered to do so by a doctor because it may cause problems

Keep out of reach of children. In case of overdose, get

medical help or contact a Poison Control Center

(1-800-222-1222, right away).

Directions

■ **do not take more than directed**

■ **the smallest effective dose should be used**

■ **do not take more than directed**

■ **adults and children 12 years and older**

■ take 1 tablet every 8 to 12 hours while symptoms last

■ for the first dose you may take 2 tablets within the first hour

■ do not exceed 2 tablets in any 8 to 12 hour period

■ do not exceed 3 tablets in a 24-hour period

■ children under 12 years: ask a doctor

Other information

■ each tablet contains: sodium 20 mg

■ store at 20-25°C (68-77°F). Avoid high humidity and

excessive heat above 40°C (104°F)

Inactive Ingredients

FD&C #2 aluminum lake, magnesium stearate,

microcrystalline cellulose, polyethylene glycol, polyvinyl

alcohol, povidone, silicon dioxide, lac, titanium dioxide

Questions or comments?

Call 1-800-540-3765

STOP PEELING

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active ingredient (in each tablet)

Naproxen sodium 220 mg

Naproxen 200 mg (NSAID)

Nonsteroidal anti-inflammatory drug

Pain Reliever/
Fever Reducer

Temporarily reduces fever

Temporarily relieves minor aches and pains due to:

■ headache

■ toothache

■ backache

■ menstrual cramps

■ minor pain of arthritis

Temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction,

especially in people allergic to aspirin. Symptoms may include:

■ asthma (wheezing)

■ skin redness

■ facial swelling

■ hives

■ rash

■ dizziness

■ loss of consciousness

■ fainting

■ difficulty breathing

■ difficulty swallowing

■ difficulty talking

■ difficulty seeing

■ difficulty hearing

■ difficulty feeling

■ difficulty moving

■ difficulty thinking

■ difficulty remembering

■ difficulty concentrating

■ difficulty understanding

■ difficulty communicating

■ difficulty interacting

■ difficulty relating

■ difficulty connecting

■ difficulty associating

■ difficulty correlating

■ difficulty contrasting

■ difficulty distinguishing

■ difficulty discerning

■ difficulty detecting

■ difficulty identifying

■ difficulty recognizing

■ difficulty knowing

■ difficulty understanding

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if

■ the stomach bleeding warning applies to you

■ you have a history of stomach problems, such as heartburn

■ you have high blood pressure, heart disease, liver cirrhosis,

kidney disease, asthma, or had a stroke

■ you are taking a

diuretic

■ you have problems or serious side effects from

taking pain relievers or fever reducers

■ This product is not manufactured or distributed by the

owner of the registered trademark Aleve®.

DIST. BY GERI-CARE PHARMACEUTICALS CORP.

1233 Vernon Ave., Cincinnati, OH 45240

REV 1224L



3 57896 95401 6

PEEL HERE FOR MORE DRUG FACTS