

Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID-nonsteroidal anti-inflammatory drug).....Pain reliever

Purpose

- Uses**
- for the temporary relief of minor aches and pains
 - ask your doctor about other uses for aspirin

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include:

- hives • facial swelling • asthma (wheezing) • shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

- Do not use**
- if you are allergic to aspirin or any other pain reliever/fever reducer
 - if you ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if • the stomach bleeding warning applies to you • you are taking a diuretic • you have a history of stomach problems, such as heartburn • you have: ◦ high blood pressure ◦ heart disease ◦ liver cirrhosis ◦ kidney disease ◦ asthma

Ask a doctor or pharmacist before use if you are • taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if • you experience any of the following signs of stomach bleeding: ◦ feel faint ◦ vomit blood ◦ have bloody or black stools ◦ have stomach pain that does not get better • an allergic reaction occurs. Seek medical help right away. • pain gets worse or lasts more than 10 days • redness or swelling is present • fever gets worse or lasts more than 3 days • any new symptoms occur • ringing in the ears or loss of hearing occurs. These may be signs of a serious condition. ➤

NDC 57896-981-10

GERI-CARE®

ADULT LOW DOSE
ENTERIC COATED

ASPIRIN 81 mg

PAIN RELIEVER (NSAID)



Compare to active ingredient
in **BAYER®** Low Dose Aspirin*

1000 Tablets

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

- Directions**
- drink a full glass of water with each dose
 - swallow whole, do not chew or crush
 - do not exceed recommended dose

adults and children 12 years and older	take 4-8 tablets every 4 hours, as needed, not more than 48 tablets in 24 hours, or as directed by a doctor
children under 12	ask a doctor

Other information • store at 20°-25°C (68°-77°F) ; excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients: cellulose, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, PEG, polydextrose, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, starch, talc titanium dioxide, triacetin, triethyl citrate, wax.

Questions or comments? 1-800-540-3765

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

*This product is not manufactured or distributed by the owner of the registered trademark Bayer®.

DIST. BY: **GERI-CARE
PHARMACEUTICALS CORP.**
1295 Towbin Avenue
Lakewood, NJ 08701



REV 0724L