

ADULT LOW DOSE  
ENTERIC COATED  
**ASPIRIN 81 mg**  
PAIN RELIEVER (NSAID)



Compare to active ingredient  
in BAYER® Low Dose Aspirin\*

120 Tablets

**Drug Facts** (continued)

**Ask a doctor before use if**  you have stomach bleeding warning applies to you  you are taking a diuretic  you have a history of stomach problems, such as heartburn  you have a high blood pressure  heart disease  liver cirrhosis  kidney disease  asthma

**Ask a doctor or pharmacist before use if you are**  taking a prescription drug for diabetes, gout or arthritis

**Stop use and ask a doctor if**  you experience any of the following signs of stomach bleeding  feel faint  vomit blood  have bloody or black stools  have stomach pain that does not get better  an allergic reaction occurs. Seek medical help right away.  pain gets worse or lasts more than 10 days  redness or swelling is present  fever gets worse or lasts more than 3 days  any new symptoms occur  ringing in the ears or loss of hearing occurs  There can be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  do not exceed recommended dose  drink a full glass of water with each dose  swallow whole, do not chew or crush

adults and children  take 4-8 tablets every 4 hours, as needed, not more than 48 tablets in 24 hours, or as directed by a doctor  
12 years and older   
  
children under 12  ask a doctor

**Other information**  store at 20-25°C (68-77°F); excursions permitted between 15°C -30°C (59°F-86°F)

**Inactive ingredients:** cellulose, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, PEG, polyacrylonitrile, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, starch, talc, titanium dioxide, triacetin, triethyl citrate, wax.

**Questions or comments?** 1-800-540-3765

STOP PEELING

**Tamper Evident:** Do not use if imprinted seal under cap is missing or broken.

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 81 mg (NSAID) ..... Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Uses**  for the temporary relief of minor aches and pains  ask your doctor about other uses for aspirin

**Warnings**

**Reye's Syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy Alert:** Aspirin may cause a severe allergic reaction which may include:  hives  facial swelling

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  are age 60 or older  have had stomach ulcers or bleeding problems  take a blood thinning (anticoagulant) or steroid drug  take other drugs containing naproxen or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  have 3 or more alcoholic drinks every day while using this product.  take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

\*This product is not manufactured or distributed by the owner of the registered trademark Bayer®.

Distributed by:

GERH-CARE PHARMACEUTICALS CORP.  
1295 Towbin Ave. Lakewood, NJ 08701

REV0322L



3 57896 98112 9

PEEL HERE FOR MORE DRUG FACTS