DKOPS GAS RELIEF <u>INFANTS'</u>

SAVE CARTON FOR COMPLETE DRUG FACTS

Drug Facts

Active ingredient (in each 0.3 mL) Simethicone 20 mg

Purpose Antigas

Use

relieves the symptoms referred to as gas

Warnings

Keep out of reach of children. In case of overdose, get medical help or contact a poison control center right away.

Directions

- shake well before using
- all dosages may be repeated as needed, after meals and at bedtime, or as directed by a physician
- do not exceed 12 doses per day
- fill enclosed dropper to recommended dosage level and dispense liquid slowly into baby's mouth, toward the inner cheek
- dosage can also be mixed with 1 oz. of cool water, infant formula or other suitable liquids
- clean dropper well after each use and replace original cap on bottle

Age (years)	Weight (lbs)	Dose
infants (under 2)	under 24	0.3 mL
children (2 and over)	24 and over	0.6 mL

Other information

- TAMPER-EVIDENT: Do not use if foil seal under the cap is broken or punctured.
- store at room temperature
- do not freeze
- see bottom panel for lot number and expiration date

Drug Facts (continued)

Inactive ingredients

carboxymethylcellulose sodium, citric acid, flavor, maltitol, microcrystalline cellulose, purified water, sodium benzoate, sodium

*This product is not manufactured

or distributed by the owner of the registered trademark MYLICON®.

citrate, xanthan gum Questions or comments?

1-800-540-3765



NDC 57896-798-01

GAS RELIEF DROPS DYE FREE

Simethicone Oral Suspension USP



Antigas Relieves the Symptoms Referred to as Gas

Plastic Dropper Enclosed

1 FL OZ (30 mL)



Compare to active ingredient of Infants MYLICON®*



Children 2 years and over

Infants under 2 years — 0.3 mL

___ 0.6 mL

Distributed by Geri-Care Pharmaceuticals 1650 63rd Street

REV 0414 410-26119-0



6.5314"