



THE SENSIBLE CHOICE
FOR PRICE AND HEALTH

NDC 57896-941-20

IBUPROFEN

PAIN RELIEVER / FEVER REDUCER (NSAID)



COMPARE TO THE ACTIVE
INGREDIENT IN ADVIL[®]

**200 Tablets
200 mg Each**

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

See carton for complete Drug Facts

Active ingredient (in each brown tablet)

Purpose

Ibuprofen USP, 200 mg (NSAID)*
*nonsteroidal anti-inflammatory drug

Pain reliever/

Fever reducer

Uses ■ Temporarily relieves minor aches and pains due to:
☐ headache ☐ toothache ☐ backache
☐ menstrual cramps ☐ the common cold
☐ muscular aches ☐ minor pain of arthritis
 ■ Temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 ■ shock ■ facial swelling ■ asthma (wheezing) ■ hives ■ rash
 ■ blisters ■ skin reddening, if an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma ■ you are taking a diuretic

*This product is not manufactured or distributed by the owner of the registered trademark Advil[®].

Distributed By: Geri-Care Pharmaceuticals Corp.

1650 63rd St., Brooklyn, NY 11204

REV 1112



3 57896 94120 8

PEEL HERE FOR MORE DRUG FACTS

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

■ under a doctor's care for any serious condition
 ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin

■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ heartburn

■ have bloody or black stools ■ vomit blood
 ■ have stomach pain that does not get better
 ■ pain gets worse or lasts for more than 10 days
 ■ fever gets worse or lasts for more than 3 days
 ■ redness or swelling is present in the painful area
 ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away.

Directions

■ **do not take more than directed**
 ■ the smallest effective dose should be used
 ■ do not take longer than 10 days, unless directed by a doctor (see Warnings)

■ adults and children 12 years and over:

■ take 1 tablet every 4 to 6 hours while symptoms persist
 ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not take more than 6 tablets in 24 hours unless directed by a doctor
 ■ children under 12 years: ask a doctor

Other information ■ store at controlled room temperature • avoid excessive heat 40°C (104°F)

Inactive ingredients: See carton

Questions or comments? Call 1-800-5-40-3765

STOP PEEING