

Drug Facts

Active ingredient (per 5mL teaspoonful) Purpose

Acetaminophen 160 mg.....Pain Reliever / Fever Reducer

Uses

• temporarily relieves minor aches and pains due to:
• headache • muscular aches • backache • minor pain of arthritis
• the common cold • toothache • premenstrual and menstrual cramps • temporarily reduces fever

Warnings

Liver Warning: This product contains acetaminophen. Severe liver damage may occur if: • adult takes more than 6 doses in 24 hours, which is the maximum daily amount • child takes more than 5 doses in 24 hours, which is the maximum daily amount • taken with other drugs containing acetaminophen • adult has 3 or more alcoholic drinks everyday while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have a liver disease.

Ask a doctor before use if you are taking the blood thinning drug warfarin.

When using this product: Do not exceed recommended dose.

Stop use and ask a doctor if • pain gets worse or lasts more than 10 days in adults • pain gets worse or lasts more than 5 days in children under 12 years • fever gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present. These could be signs of a serious condition.

NDC 57896-160-16



LIQUID PAIN RELIEF

ACETAMINOPHEN
160 mg / 5 mL
PAIN RELIEVER • FEVER REDUCER

ASPIRIN FREE • ALCOHOL FREE
CHERRY FLAVOR

Compare to the active ingredient of
Tylenol® Liquid.*

THIS BULK PACKAGE IS NOT FOR HOUSEHOLD USE.
Dispense contents with a child resistant closure in a tight, light-resistant container as defined in the USP.

*This product is not manufactured or distributed by the owner of the registered trademark Tylenol®.

16 FL OZ (473 mL)

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults/children even if you do not notice any signs or symptoms.

Directions • do not take more than directed

- find dose in chart below
- if possible, use weight to dose, otherwise use age
- dosage may be repeated every 4 hours, or as directed by a doctor
- do not give more than 5 doses in 24 hours

Weight (lbs)	under 24	24-35	36-47	48-59	60-71	72-95	over 95
Age (years)	Under 2	2 to 3	4 to 5	6 to 8	9 to 10	11	12 to adult
Dosage Teaspoonful (tsp)	consult Physician	1 tsp (5 mL)	1 ½ tsp (7.5 mL)	2 tsp (10 mL)	2 ½ tsp (12.5 mL)	3 tsp (15 mL)	4 tsp (20 mL)

Other information

- for institutional or pharmacy use only
- store at room temperature 15°- 30°C (59°-86°F)
- protect from freezing • protect from light
- **TAMPER-EVIDENT:** Do not use if foil seal over bottle opening is torn, broken or missing.

Inactive ingredients: artificial and natural cherry flavor, citric acid, FD&C Red #40, methylparaben, polyethylene glycol, propylene glycol, propylparaben, purified water, sodium citrate, sucralose

Code # L150

DIST. BY:

Geri-Care Pharmaceuticals Corp.

1650 63rd Street

Brooklyn, NY 11204

Rev.:05/19

Lot#

Exp. Date:

