

Size: 3.25 x 7"

Colors: 4CP



IRON SUPPLEMENT LIQUID

FERROUS SULFATE
220 mg/5 mL

For the treatment of iron deficiency and
iron deficiency anemia

16 fl oz (473 mL)

Supplement Facts

Serving Size: 1 teaspoonful (5 mL)

Servings per container: 94

	Amount per teaspoonful (5 mL)	%Daily Value*
Calories	4	
Total Carbohydrate	1g	<1%
Total Sugars	1g	**
Includes 1g Added Sugars		2%
Iron (elemental) (from ferrous sulfate)	44mg	244%

*% Daily Value tells you how much a nutrient in a serving of food contributes to a daily diet. 2,000 calories a day is used for general nutrition advice.

**Daily Value not established

Each teaspoonful (5mL) contains 220mg Ferrous Sulfate, providing approximately 44mg of elemental iron.

OTHER INGREDIENTS: Purified water, sucrose, artificial and natural cherry flavor, citric acid, sodium benzoate, sucralose, FD&C yellow #6.

WARNING: Accidental overdose of iron containing supplements is a leading cause of fatal poisoning in children under 6.

KEEP THIS PRODUCT OUT OF REACH OF CHILDREN.
In case of accidental overdose, call a doctor or
Poison Control Center immediately.

Distributed by:
GERI-CARE Pharmaceuticals Corp.
1650 63rd Street, Brooklyn, NY 11204

SUGGESTED USE: Adults and children 12 years and over, take one (1) teaspoonful daily with or after a meal as a dietary supplement, or as directed by your doctor. Children under 12, consult a doctor.

Mix with water or juice to avoid temporary staining of teeth. Do not mix with milk or wine-based vehicles.

CAUTION: Do not exceed suggested serving size. The treatment of any anemic condition should be under the advice and supervision of a doctor. Since oral iron products interfere with absorption of oral tetracycline antibiotics, these products should not be taken within two (2) hours of each other.

This product may occasionally cause constipation, diarrhea, or may color stool black. It may cause gastrointestinal discomfort (such as nausea) which may be minimized by taking with meals.

As with any supplement, if you are pregnant, nursing, taking medication, or under a doctor's care for any condition, ask a health professional before use.

OTHER INFORMATION: Store at room temperature 15-30°C (59-86°F)

Tamper-Evident: Do not use if foil seal over bottle opening is torn, broken or missing.

Pharmacist: Dispense in a tight, light-resistant container with a child resistant cap as defined in the USP

FOR INSTITUTIONAL USE ONLY

Code#L-57
Rev.:0719

Lot #
Exp. Date:

