

Drug Facts

Active ingredient Sorbitol 70%, USP.....
Purpose Laxative

Uses

- relieves occasional constipation and irregularity
- generally produces bowel movement in ¼ to 1 hour when used rectally
- as a pharmaceutical aide (sweetener)
- for other uses, ask your doctor

Warnings

For rectal use only

- Do not use** • when abdominal pain, nausea, or vomiting are present
- for more than one week unless directed by a doctor.

Ask a doctor before use if you

- are taking mineral oil
- have noticed a sudden change in bowel habits that lasts over 2 weeks

Stop use and ask a doctor if you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. →

Code#: L-126

Rev. 12/20

Lot.#

Exp. Date:

NDC 57896-758-16

GERI-CARE®

SORBITOL SOLUTION USP 70% W/W LAXATIVE

TAMPER EVIDENT: Do not use this product if inner foil seal over the mouth of bottle is missing or broken.

16 FL OZ (1 PT) (473 mL)

Drug Facts (continued)

Directions • do not exceed recommended dose

Age	Dose
Adults and children 12 years and over	For rectal use only. Enema dosage is 120 mL of a 25 to 30% w/v solution (1 part of this product with 2.3 parts water) in a single daily dose as needed, or as directed by your doctor.
Children under 12 years of age	Ask a doctor.

Other information

- store at room temperature 15°- 30° C (59°-86°F)
- below 59°F cloudiness and thickening may occur; warming will restore clarity and fluidity without affecting product quality
- do not freeze
- Pharmacist-Preserve and dispense in tight, light-resistant container with a child resistant cap as defined in the USP

Inactive ingredient: water

Questions or comments? 1-800-540-3765



**DIST. BY: GERI-CARE
PHARMACEUTICALS CORP.**
1650 63rd Street
Brooklyn, NY 11204