

SAVE CARTON FOR COMPLETE DRUG FACTS

Drug Facts

Active ingredient **Purpose**
(in each 0.3 mL)
Simethicone 20 mg Antigas

Use
relieves the symptoms referred to as gas

Warnings
Keep out of reach of children. In case of overdose, get medical help or contact a poison control center right away.

Directions

- shake well before using
- all dosages may be repeated as needed, after meals and at bedtime, or as directed by a physician
- do not exceed 12 doses per day
- fill enclosed dropper to recommended dosage level and dispense liquid slowly into baby's mouth, toward the inner cheek
- dosage can also be mixed with 1 oz. of cool water, infant formula or other suitable liquids
- clean dropper well after each use and replace original cap on bottle

Age (years)	Weight (lbs)	Dose
infants (under 2)	under 24	0.3 mL
children (2 and over)	24 and over	0.6 mL

Other information

- **TAMPER-EVIDENT:** Do not use if foil seal under the cap is broken or punctured.
- store at room temperature
- do not freeze
- see bottom panel for lot number and expiration date

Drug Facts
(continued)

Inactive ingredients
carboxymethylcellulose
sodium, citric acid, flavor,
maltitol, microcrystalline
cellulose, purified water,
sodium benzoate, sodium
citrate, xanthan gum

Questions or comments?
1-800-540-3765

*This product is not manufactured or distributed by the owner of the registered trademark MYLICON®.

INFANTS' GAS RELIEF DROPS



NDC 57896-798-01



THE SENSIBLE CHOICE
FOR PRICE AND HEALTH

INFANTS' GAS RELIEF DROPS **DYE FREE**

Simethicone Oral Suspension USP



Antigas
Relieves the Symptoms
Referred to as Gas

Plastic Dropper Enclosed

1 FL OZ
(30 mL)



Compare to active
ingredient of
Infants MYLICON®



DOSAGE

Children
2 years
and over

0.6 mL

Infants
under
2 years

0.3 mL

DROPPER ENCLOSED

Distributed by
Geri-Care Pharmaceuticals
1650 63rd Street
Brooklyn, NY 11204

REV 0414
410-26119-0



N 3 57896-79801 7