

<p><b>Drug Facts (continued)</b></p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>■ If you have ever had an allergic reaction to any other pain reliever or a fever reducer</li> <li>■ For strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.</li> <li>■ For the temporary relief of arthritis pain ONLY in the following areas: <ul style="list-style-type: none"> <li>■ hand, wrist, elbow (upper body areas)</li> <li>■ foot, ankle, knee (lower body areas)</li> </ul> </li> <li>■ This product may take up to 7 days to work for arthritis pain; it is not for immediate relief.</li> <li>■ If no pain relief in 7 days, stop use.</li> </ul> <p><b>Warnings</b></p> <p><b>For external use only</b></p> <p><b>Allergy alert:</b> Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: <ul style="list-style-type: none"> <li>■ hives</li> <li>■ asthma (wheezing)</li> <li>■ skin redness</li> <li>■ blisters</li> <li>■ facial swelling</li> <li>■ shock</li> <li>■ rash</li> </ul> If an allergic reaction occurs, stop use and seek medical help right away.</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Ask a doctor or pharmacist before use if you are</b></p> <ul style="list-style-type: none"> <li>■ taking any other drug</li> <li>■ under a doctor's care for any serious condition</li> <li>■ under age 18 years. It is not known if this drug works or is safe in children</li> </ul> <p><b>Ask a doctor before use if</b></p> <ul style="list-style-type: none"> <li>■ you have problems or serious side effects from taking pain relievers or fever reducers</li> <li>■ stomach bleeding warning applies to you</li> <li>■ you have a history of stomach problems, such as heartburn</li> <li>■ you have high blood pressure, heart disease, kidney disease, asthma, or had a stroke</li> <li>■ you are taking a diuretic</li> <li>■ you are under age 18 years.</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>■ skin irritation occurs</li> <li>■ fever occurs</li> <li>■ pain gets worse or lasts more than 21 days</li> <li>■ redness or swelling is present in the painful area</li> <li>■ you experience any of the following signs of a serious condition: <ul style="list-style-type: none"> <li>■ any new symptoms appear. These could be signs of stomach bleeding.</li> <li>■ you have bloody or black stools</li> <li>■ have stomach or black stools</li> <li>■ feel faint</li> <li>■ vomit blood</li> <li>■ have symptoms of heart problems or stroke: <ul style="list-style-type: none"> <li>■ chest pain</li> <li>■ trouble breathing</li> <li>■ leg swelling</li> <li>■ slurred speech</li> </ul> </li> <li>■ weakness in one part or side of body</li> </ul> </li> </ul>
--	---

<p><b>Drug Facts (continued)</b></p> <p><b>Liver warning:</b> This product contains diclofenac. Liver damage may occur if you apply when using other drugs containing diclofenac</p> <p><b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is small but higher if you</p> <ul style="list-style-type: none"> <li>■ are age 60 or older</li> <li>■ have had stomach ulcers or bleeding problems</li> <li>■ take a blood thinning (anticoagulant) or steroid drug</li> <li>■ take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)</li> <li>■ have 3 or more alcoholic drinks every day while using this product</li> <li>■ apply more or for longer than directed</li> </ul> <p><b>Heart attack and stroke warning:</b> NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Active ingredient</b></p> <p>Diclofenac sodium (NSAID) 1%..... Arthritis pain reliever</p> <p><b>Purpose</b></p> <p>Arthritis pain reliever</p> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>■ for the temporary relief of arthritis pain ONLY in the following areas: <ul style="list-style-type: none"> <li>■ hand, wrist, elbow (upper body areas)</li> <li>■ foot, ankle, knee (lower body areas)</li> </ul> </li> <li>■ this product may take up to 7 days to work for arthritis pain; it is not for immediate relief.</li> <li>■ If no pain relief in 7 days, stop use.</li> </ul> <p><b>Warnings</b></p> <p><b>For external use only</b></p> <p><b>Allergy alert:</b> Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: <ul style="list-style-type: none"> <li>■ hives</li> <li>■ asthma (wheezing)</li> <li>■ skin redness</li> <li>■ blisters</li> <li>■ facial swelling</li> <li>■ shock</li> <li>■ rash</li> </ul> If an allergic reaction occurs, stop use and seek medical help right away.</p>
--	---

### Use Enclosed Dosing Card to Measure a Dose

**Diclofenac Sodium Topical Gel, 1% (NSAID)- arthritis pain reliever**

**Dosing Card (NOT ACTUAL SIZE)**

**Read product User Guide before use**

**Not for immediate pain relief**

Upper body dose 2.25 inches long (2 grams)

Lower body dose 4.5 inches long (4 grams)

**Tamper evident statement: Do not use if safety seal on tube nozzle is broken or missing.**

### How to Use

- For arthritis pain only
- Use 4 times a day
- May take up to 7 days to work for your arthritis pain
- For use on no more than 2 body areas
- Use up to 21 days unless directed by your doctor



NDC 57896-140-01

Compare to the active ingredient in Voltaren Arthritis Pain Gel\*\*

# Diclofenac Sodium Topical Gel, 1%

(NSAID) - Arthritis Pain Reliever

NET WT 3.53 oz (100 g)

For external use only

### Original Prescription Strength

- For Daily Treatment of Arthritis Pain
- Anti-Inflammatory

**Medicated Gel**  
**Clinically Proven to**  
**Relieve Arthritis Pain**



Hand



Wrist



Elbow



Foot



Ankle



Knee

Not for use on any other body area (such as back, hip, or shoulder)

Distributed by:  
GERI-CARE PHARMACEUTICALS CORP.  
1650 63rd Street  
Brooklyn, NY 11204

Made in India Code: GO/DRUGS/361

\*\*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, PLC, distributor of Voltaren Arthritis Pain Gel.

REV: 10/20

DDI3B/00

<p><b>Drug Facts (continued)</b></p> <p><b>Read the enclosed User Guide for complete instructions:</b></p> <ul style="list-style-type: none"> <li>■ use only as directed</li> <li>■ do not use more than directed or for longer than directed</li> <li>■ apply only to clean, dry skin that does not have any cuts, open wounds, infections or rashes</li> <li>■ do not apply in same area as any other product</li> <li>■ do not apply with external heat such as heating pad</li> <li>■ do not apply a bandage over the treated area</li> <li>■ store ENCLOSED DOSING CARD with your Diclofenac Sodium Topical Gel. The dosing card is re-usable.</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ Store at 20-25°C (68-77°F). Keep from freezing.</li> <li>■ read all product information before using. Keep the dosing card, this carton and accompanying User Guide for important information.</li> </ul> <p><b>Inactive ingredients</b> Carbomer homopolymer Type C, cocoyl caprylocaprate, isopropyl alcohol, mineral oil, polyoxy 20 cetostearyl ether, propylene glycol, purified water, and strong ammonia solution.</p> <p><b>Questions or comments? 1-888-375-3784</b></p>	<p><b>Drug Facts (continued)</b></p> <p><b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p> <p><b>Directions</b></p> <p>Use up to 21 days unless directed by your doctor.</p> <p>Not for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.</p> <p><b>For your arthritis pain:</b></p> <ul style="list-style-type: none"> <li>■ Use 4 times per day every day</li> <li>■ Use 4 times per body area (hand, wrist, or elbow)</li> <li>■ Squeeze out 2.25 inches (2 grams)</li> <li>■ For each lower body area (foot, ankle or knee) - Squeeze out 4.5 inches (4 grams)</li> </ul> <p><b>Per Dose</b></p> <p>Use ENCLOSED DOSING CARD to measure a dose</p>
---	---



3 57896 14001 4