

Drug Facts
Keep Outer Package for Complete Product Information
Active Ingredient (in each caplet) Loperamide HCl 2 mg
Purpose Anti-diarrheal
 Use controls symptoms of diarrhea, including Traveler's Diarrhea

NDC 57896-381-13



ANTI-DIARRHEAL

Loperamide HCl Tablets, 2 mg

Anti-Diarrheal

Controls the Symptoms of Diarrhea



compare to active ingredient in Imodium® A-D*

12 caplets

TAMPER EVIDENT DO NOT USE IF PACKAGE IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING
 * This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark, Imodium® A-D. 50844 OR6091937502
 REV. 03/21
DISTRIBUTED BY:
GERI-CARE PHARMACEUTICAL CORP.
 1650 63 Street, Brooklyn, NY 11204



- To open, tear at perforations and remove a section.
- Use to eat from caplet.

LOT/EXP ONLY

Drug Facts (continued)
Questions or comments? Call 1-800-540-3765

3 57896 38113 4

Drug Facts (continued)
Other Information ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN ■ Store between 20°-25°C (68°-77°F) ■ See end flap for expiration date and lot number
Inactive ingredients corn starch, D&C yellow #10 aluminum lake, magnesium stearate, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, microcrystalline cellulose, dibasic calcium phosphate dihydrate, silicon dioxide

Drug Facts (continued)
Warnings
Heart alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.
Heart alert: Taking more than directed can cause serious heart problems or death.
Do not use if you have bloody or black stool.
Ask a doctor before use if you have ■ a fever ■ mucus in the stool ■ a history of liver disease ■ a history of abnormal heart rhythm
Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.
When using this product, tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.
Stop use and ask a doctor if ■ symptoms get worse ■ diarrhea lasts for more than 2 days ■ you get abdominal swelling or bloating. These may be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions ■ Drink plenty of clear fluids to help prevent dehydration caused by diarrhea ■ Find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

B-0617-375-02
 086061937502

C7354 LEBG81 6E