

Drug Facts (continued)	Drug Facts
When using this product • avoid contact with the eyes Stop use and ask a doctor if • sore mouth symptoms do	Purpose ing the ing the second
not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash or fever develops	Uses Uses pain associated with the following mouth
Keep out of the reach of children. In case of overdose, get medical help or contact a Poison	and gum irritations: • toothache • sore gums • canker sores • braces • minor dental procedures • dentures
Control Center right away. Directions	spnimeW
qo uot exceed recommended dosage	: DNINRAW AIMENIBOJOOMEHTEM
• do not use for more than 7 days unless told to do so by a	Use of this product may cause methemoglobinemia, a
<ul> <li>edutist or doctor</li> <li>adults and children 2 years of age or older: apply to</li> </ul>	serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can
attected area up to 4 times daily or as directed by	occur even if you have used this product before. Stop use
doctor/dentist	and seek immediate medical attention if you or a child in
<ul> <li>children between 2 and 12 years of age: adult</li> </ul>	your care develops:
<ul> <li>supervision should be given in use of this product</li> <li>children under 2 years of age: do not use</li> </ul>	<ul> <li>pale; gray, or blue colored skin (cyanosis)</li> </ul>
	<ul> <li>headache</li> <li>rapid heart rate</li> </ul>
Other information	<ul> <li>shortness of breath</li> </ul>
<ul> <li>do not use tube if it is cut prior to opening</li></ul>	<ul> <li>dizziness or lightheadedness</li> </ul>
<ul> <li>cut open tip of tube on score mark with scissors</li> <li>store at a controlled room temperature 68°-77°F (20°-25°C)</li> </ul>	<ul> <li>fatigue or lack of energy</li> </ul>
	Allergy alert: do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine,
<b>Inactive ingredients</b> Flavor, PEG-6, PEG-8, PEG-32, PEG-75, Sodium Saccharin,	benzocaine, or other "caine" anesthetics. If a skin reaction
Sorbic Acid	occurs, stop use and seek medical help right away.
<b>Questions or Comments?</b> 1-800-540-3765	Do not use • for teething • in children under 2 years of age
	<ul> <li>more than recommended</li> <li>for more than 7 days unless told to do so by a dentist or doctor</li> </ul>

Distributed by: Geri-Care Pharmaceuticals 1650 63rd St. Brooklyn NY 11204 **REV 1020** 

\*This product is not manufactured or distributed by © CHURCH & DWIGHT CO., INC., owner of the registered trademark Orajel®

**Instant Oral Pain Relief** 

**OOTHACHE &** 

**Temporary Pain Relief for:** Occasional minor irritation

INSTANT

 Sore mouth Toothache

Ę

ГОТ

20% BENZOCAINE TOPICAL GEL

NET WT. 0.5 oz. (14.2g)

13905<sup>III</sup>

6


57896

3

NDC 57896-139-05



Tube safety sealed for your protection



**RAL PAIN RELIEF** 

Compare to active ingredient in Orajel <sup>®</sup> MAXIMUM Toothache and Oral Pain Relief