



COATING
FREE AREA



1 7 8 9 9 9



Drug Facts

Active ingredient (in each suppository)

Bisacodyl USP 10 mg.....Stimulant laxative

Purpose

Uses

- relieves occasional constipation
- this product generally produces a bowel movement in 15 minutes to 1 hour

Warnings

For rectal use only

Do not use when abdominal pain, nausea, or vomiting are present.

Ask a doctor before use if you have a sudden change in bowel habits that lasts over a period of 2 weeks.

When using this product you may experience

- rectal burning ■ mild cramps ■ abdominal discomfort ■ faintness

Stop use and ask a doctor if ■ you have rectal bleeding or you fail to have a bowel movement after using a laxative. These may be signs of a serious condition.


- you need to use a laxative for more than one week.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact Poison Control Center right away.

Directions

- tear off one suppository from the strip
- peel apart the wrapper
- continue peeling apart to almost the full length of the suppository
(see illustration at the right)
- gently remove the suppository from the wrapper
- lie on your side and push suppository, with pointed end first, high into the rectum so it will not slip out
- retain it for 15 to 20 minutes
- **Do not exceed recommended dose**
- if suppository seems soft, place in refrigerator for a short time before use



Adults and children 12 years and over	1 suppository once daily
Children under 12 years	Ask a doctor

*This product is not manufactured or distributed by the owner of the registered trademark Dulcolax®. Distributed by: Qualitabs Inc. Brooklyn, NY 11204
 Questions or Comments: 1-800-540-3765



REV1119

Drug Facts (continued)
Other information ■ store at temperatures below 25°C (77°F) ■ Made in India.
Inactive ingredient hard fat

Compare to the active ingredient in Dulcolax® Suppositories*

NDC 57896-443-01



Bisacodyl Suppositories

Fast Acting Stimulant Laxative

100 Suppositories

10 mg each

TAMPER EVIDENT: SUPPOSITORES ARE WRAPPED IN TAMPER EVIDENT SEALED WRAPPERS. DO NOT USE IF WRAPPER IS TORN OR OPEN WHEN PURCHASED.



Bisacodyl Suppositories