

HEALTH  STAR®

NDC 57896-911-36

ADULT LOW DOSE
CHEWABLE

ASPIRIN 81 mg
PAIN RELIEVER (NSAID)



Compare to active ingredient
in **BAYER®** Low Dose Aspirin*

36 Tablets

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses ■ for the temporary relief of minor aches and pains
■ ask your doctor about other uses for aspirin

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling
■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if ■ the stomach bleeding warning applies to you ■ you are taking a diuretic ■ you have a history of stomach problems, such as heartburn
■ you had a tonsillectomy or oral surgery within the past 7 days

■ you have: high blood pressure heart disease
 liver cirrhosis kidney disease asthma

* This product is not manufactured or distributed by the owner of the registered trademark Bayer®.

Dist. by: **Gerl-Care Pharmaceuticals Corp.**
1650 63rd Street, Brooklyn, NY 11204

REV 0421



PEEL HERE FOR MORE DRUG FACTS

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- taking any other drug containing an NSAID (prescription or nonprescription) ■ taking a blood thinning (anticoagulant) or steroid drug ■ taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding:

- feel faint vomit blood have bloody or black stools have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away. ■ pain gets worse or lasts more than 10 days
- redness or swelling is present ■ fever gets worse or lasts more than 3 days ■ any new symptoms occur
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- do not exceed recommended dose

adults and children 12 years and older	take 4-8 tablets every 4 hours, as needed, not more than 48 tablets in 24 hours, or as directed by a doctor
children under 12	ask a doctor

Other information ■ store at room temperature 15°C - 30°C (59°F - 86°F)

Inactive ingredients: dextrates, FD&C yellow #6, flavor, saccharin sodium, starch.

Questions or comments? 1-800-540-3765

STOP PEELING