

HEALTH STAR[®]

IBUPROFEN

200 mg

PAIN RELIEVER/ FEVER REDUCER
(NSAID)Compare to the active
ingredient in **ADVIL[®]***

50 Tablets

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active ingredient (in each brown tablet) *Paracetamol USP, 200 mg (NSAID) *... Pain reliever/fever reducer/ nonsteroidal anti-inflammatory drug*

Uses ■ temporarily relieves minor aches and pains due to:
 □ headache □ muscular aches □ toothache
 □ backache □ menstrual cramps □ the common cold
 □ minor pain of arthritis ■ temporarily reduces fever

Warnings

Always alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 ■ shock ■ facial swelling ■ asthma (wheezing) ■ skin redness
 ■ hives ■ rash ■ dizziness ■ skin redness
 use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ take more or for a longer time than directed ■ are taking aspirin, aspirin-containing products or steroids ■ are age 60 or older ■ take other drugs (including prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have had stomach ulcers or bleeding problems ■ have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not take ibuprofen if you are or may become pregnant. Do not take ibuprofen if you have ever had an allergic reaction to any other pain reliever/fever reducer.

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■

*This product is not manufactured or distributed by the owner of the registered trademark Advil.
 Distributed by: GERRI-CARE PHARMACEUTICALS CORP.
 1235 Towbin Avenue, Lakewood, NJ 08701



REV 0724

3 57896 94105 5

PEEL HERE FOR MORE DRUG FACTS

Spot UV

Drug Facts (continued)

■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are
 ■ taking aspirin, aspirin-containing products, or other drugs for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
 ■ taking any other drug

When using this product

■ take with food or milk if stomach upset occurs
Stop use and ask a doctor if you experience any of the following signs of stomach bleeding:
 ■ black or bloody stools ■ vomit blood
 ■ have stomach pain that does not get better

■ you have symptoms of heart problems or stroke
 ■ chest pain □ trouble breathing □ leg swelling
 ■ slurred speech □ weakness in one part or side of body
 ■ pain gets worse or lasts for more than 10 days
 ■ fever gets worse or lasts for more than 3 days
 ■ redness or swelling is present in the painful area
 ■ any new symptoms appear

(If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions **do not take more than directed**
 ■ the **smallest effective dose should be used**

■ adults and children 12 years and over:
 ■ take 1 tablet every 4 to 6 hours while symptoms persist
 ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not take more than 6 tablets in 24 hours unless directed by a doctor
 ■ children under 12 years: ask a doctor

Other information ■ store between 20°-25°C (68°-77°F) ■ avoid excessive heat 40°C (104°F)

Inactive ingredients: carmellose, cellulose, colloidal silicon dioxide, corn starch, hypromellose, lactose, magnesium stearate, polydextrose, PEG, iron oxide, sodium starch glycolate, stearic acid, titanium dioxide.

Questions or comments? Call 1-800-540-3765

STOP PEELING