

GERI CARE®

ADULT LOW DOSE
ENTERIC COATED
ASPIRIN 81 mg
PAIN RELIEVER (NSAID)

Compare to active ingredient
in **BAYER®** Low Dose Aspirin*

200 Tablets

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active ingredient (in each tablet)
Aspirin 81 mg (NSAID).....Pain reliever
*nonsteroidal anti-inflammatory drug

Uses ■ for the temporary relief of minor aches and pains
■ ask your doctor about other uses for aspirin

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling

■ shock ■ asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product. ■ take more or for a longer time than directed.

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if ■ the stomach bleeding warning applies to you ■ you are taking a diuretic ■ you have a history of stomach problems, such as heartburn ■ you have: □ high blood pressure □ heart disease □ liver cirrhosis □ kidney disease □ asthma

*This product is not manufactured or distributed by the owner of the registered trademark Bayer®.
Dist. by: Geri-Care Pharmaceuticals Corp.
1650 63rd St.
Brooklyn, NY 11204

REV 0721L



3 57896 98120 4

PEEL HERE FOR MORE DRUG FACTS →

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are ■ ■ ■ ■ ■
■ taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away.

■ you experience any of the following signs of stomach bleeding: □ feel faint □ vomit blood □ have bloody or black stools □ have stomach pain that does not get better

■ pain gets worse or lasts more than 10 days
■ redness or swelling is present ■ fever gets worse or lasts more than 3 days ■ any new symptoms occur
■ ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not exceed recommended dose
■ drink a full glass of water with each dose
■ swallow whole, do not chew or crush

adults and children ■ take 4-8 tablets every 4 hours, as needed, not more than 48 tablets in 24 hours, or as directed by a doctor
children under 12 ■ ask a doctor

Other information

■ store at 20-25°C (68-77°F); excursions permitted between 15°C - 30°C (59°F - 86°F)

Inactive ingredients: cellulose, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, PEG, polyethylene glycol, stearic acid, sodium bicarbonate, sodium lauryl sulfate, starch, talc, titanium dioxide, triacetin, triethyl citrate, wax.

Questions or comments? 1-800-540-3765

STOP PEELING