

Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses • for the temporary relief of minor aches and pains
• ask your doctor about other uses for aspirin

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if • the stomach bleeding warning applies to you • you are taking a diuretic • you have a history of stomach problems, such as heartburn • you have: ° high blood pressure ° heart disease ° liver cirrhosis ° kidney disease ° asthma

Ask a doctor or pharmacist before use if you are • taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if • you experience any of the following signs of stomach bleeding: ° feel faint ° vomit blood ° have bloody or black stools ° have stomach pain that does not get better • an allergic reaction occurs. Seek medical help right away. • pain gets worse or lasts more than 10 days • redness or swelling is present • fever gets worse or lasts more than 3 days • any new symptoms occur • ringing in the ears or loss of hearing occurs

NDC 57896-983-10

GERI CARE®

ADULT LOW DOSE
ENTERIC COATED

ASPIRIN 81 mg

PAIN RELIEVER (NSAID)



Compare to active ingredient
in **BAYER®** Low Dose Aspirin*

PACKAGE NOT CHILD RESISTANT
FOR INSTITUTIONAL USE ONLY

1000 Tablets

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • drink a full glass of water with each dose
• swallow whole, do not chew or crush • do not exceed recommended dose

adults and children 12 years and older	take 4-8 tablets every 4 hours, as needed, not more than 48 tablets in 24 hours, or as directed by a doctor
children under 12	ask a doctor

Other information • store at room temperature 15°C - 30°C (59°F - 86°F)

Inactive ingredients: cellulose, croscarmellose sodium, D&C Yellow #10 Lake, lactose monohydrate, methacrylic acid co-polymer, PEG, polacrillin potassium, polyvinyl alcohol, silica, sodium lauryl sulfate, soya powder, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-540-3765

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

*This product is not manufactured or distributed by the owner of the registered trademark Bayer®.

Product of India
DIST. BY: GERI-CARE
PHARMACEUTICALS CORP.
1650 63rd Street
Brooklyn, NY 11204

REV 0321B

