

ADULT LOW DOSE
ENTERIC COATED
ASPIRIN 81 mg
PAIN RELIEVER (NSAID)

Compare to active ingredient
in **BAYER[®] Low Dose Aspirin[®]**

120 Tablets

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active ingredient (in each tablet) *Aspirin 81 mg (NSAID)** Pain reliever
*nonsteroidal anti-inflammatory drug

Uses ■ for the temporary relief of minor aches and pains
■ ask your doctor about other uses for aspirin

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing aspirin or other NSAIDs (aspirin, ibuprofen, naproxen, etc.) ■ take 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

*This product is not manufactured or distributed by the owner of the registered trademark Bayer[®].

Product of India

Dist. by: Gen-Care Pharmaceuticals Corp.
1650 63rd Street, Brooklyn, NY 11204

REV 1121B



3 57896198112 9

PEEL HERE FOR MORE DRUG FACTS

Spot UV

Drug Facts (continued)

Ask a doctor before use if ■ the stomach bleeding warning applies to you ■ you are taking a diuretic ■ you have a history of stomach problems, such as heartburn ■ you have a high blood pressure ■ heart disease ■ liver cirrhosis ■ kidney disease ■ asthma

Ask a doctor or pharmacist before use if you are ■ taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ an allergic reaction occurs. Seek medical help right away ■ pain gets worse or lasts more than 10 days ■ redness or swelling is present ■ fever gets worse or lasts more than 3 days ■ any new symptoms occur ■ ringing in the ears or loss of hearing occurs. These can be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not exceed recommended dose ■ drink a full glass of water with each dose ■ swallow whole; do not chew or crush

adults and children take 4-8 tablets every 4 hours, as needed; not more than 48 tablets in 24 hours, or as directed by a doctor

children under 12 ask a doctor
Other information ■ store at controlled room temperature between 20°-25°C (68°-77°F)

Inactive ingredients: cellulose, croscarmellose sodium, D&C Yellow #10 Lake, lactose monohydrate, methacrylic acid co-polymer, PEG, polacrillin potassium, polyvinyl alcohol, silica, sodium lauryl sulfate, soya powder, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-540-3765

STOP PEELING