

GERI CARE

NDC 57896-941-50

IBUPROFEN

PAIN RELIEVER / FEVER REDUCER

(NSAID)

COMPARE TO THE ACTIVE
INGREDIENT IN ADVIL®*

PACKAGE NOT CHILD RESISTANT

500 Tablets
200 mg Each

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active ingredient (in each brown tablet)

ibuprofen USP, 200 mg (NSAID)*Pain reliever / fever reducer
*nonsteroidal anti-inflammatory drug

Uses • Temporarily relieves minor aches and pains due to: • headaches • menstrual cramps • muscular aches • toothache • the common cold • minor pain of arthritis • backache • Temporarily reduces fever

Warnings

Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- shock
 - facial swelling
 - asthma (wheezing)
 - hives
 - rash
 - skin reddening
 - blisters
- if an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

Do not use • if you have ever had an allergic reaction to any other pain reliever/fever reducer • right before or after heart surgery

Ask a doctor before use if • stomach bleeding warning applies to you • you have problems or serious side effects from taking pain relievers or fever reducers • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma • you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product • take with food or milk if stomach upset occurs • the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear
- if pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • do not take more than directed • the smallest effective dose should be used • do not take longer than 10 days, unless directed by a doctor (see Warnings) • adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist • if pain or fever does not respond to 1 tablet, 2 tablets may be used • do not take more than 6 tablets in 24 hours unless directed by a doctor • children under 12 years: ask a doctor

Other information • store between 20°-25°C (68°-77°F) • avoid excessive heat 40°C (104°F)

Inactive ingredients: carnauba wax, cellulose, corn starch, fumed silica gel, hypromellose, lactose, magnesium stearate, polydextrose, PEG, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide.

For institutional use only.

REV 0213

*This product is not manufactured or distributed by the owner of the registered trademark Advil®. call 1-800-540-3765

Questions or comments?



3 57896 94150 5

Dist. by: **Gert-Care**
Pharmaceuticals Corp.
1650 63rd Street
Brooklyn, NY 11204