

VALUE SIZE NDC 57896-981-20

GERI-CARE®

**Adult
Low Dose
Enteric Coated
Aspirin
Pain Reliever (NSAID)**

COMPARE
TO ACTIVE
INGREDIENT
IN BAYER®
LOW DOSE
ASPIRIN*

PACKAGE NOT CHILD RESISTANT

200 Tablets 81 mg each

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)* Pain reliever
*nonsteroidal anti-inflammatory drug

Uses ■ for the temporary relief of minor aches and pains
■ ask your doctor about other uses for aspirin

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling

■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if ■ the stomach bleeding warning applies to you ■ you are taking a diuretic ■ you have a history of stomach problems, such as heartburn ■ you have: high blood pressure heart disease liver cirrhosis kidney disease asthma

*This product is not manufactured or distributed by the owner of the registered trademark Bayer®.

Distributed By: **Geri-Care Pharmaceuticals Corp.**
1650 63rd Street, Brooklyn, NY 11204



REV 1018

3 57896 98120 4

PEEL HERE FOR MORE DRUG FACTS

Drug Facts(continued)

Ask a doctor or pharmacist before use if you are

- taking any other drug containing an NSAID (prescription or nonprescription) ■ taking a blood thinning (anticoagulant) or steroid drug ■ taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if ■ you experience any

of the following signs of stomach bleeding:

- feel faint □ vomit blood □ have bloody or black stools □ have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away. ■ pain gets worse or lasts more than 10 days ■ redness or swelling is present
- fever gets worse or lasts more than 3 days
- any new symptoms occur ■ ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health

professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- swallow whole, do not chew or crush
- do not exceed recommended dose

adults and children
12 years and older

take 4-8 tablets every 4 hours,
as needed, not more than
48 tablets in 24 hours, or as
directed by a doctor

children under 12

ask a doctor

Other information ■ store at controlled room temperature

Inactive ingredients: cellulose, D&C yellow

#10 lake, FD&C yellow #6 lake, hypromellose, methacrylic acid, PEG, polydextrose, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, starch, talc, titanium dioxide, triacetin, triethyl citrate, wax.

STOP PEELING