



NDC 57896-981-30

THE SENSIBLE CHOICE FOR PRICE AND HEALTH

# ADULT LOW DOSE ENTERIC COATED ASPIRIN

PAIN RELIEVER (NSAID)

COMPARE TO THE ACTIVE INGREDIENT IN BAYER® LOW DOSE ASPIRIN\*

300 Tablets  
81 mg each

**Tamper Evident:** Do not use if imprinted seal under cap is missing or broken.

## Drug Facts (see carton for complete Drug Facts).

**Active ingredient (in each tablet)** **Purpose**  
Aspirin 81 mg (NSAID)\* ..... Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Uses** ■ for the temporary relief of minor aches and pains  
■ ask your doctor about other uses for aspirin

## Warnings

**Reye's Syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy Alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use** if you are allergic to aspirin or any other pain reliever/fever reducer.

**Ask a doctor before use if** ■ the stomach bleeding warning applies to you ■ you are taking a diuretic ■ you have a history of stomach problems, such as heartburn ■ you have:  high blood pressure  heart disease  liver cirrhosis  kidney disease  asthma

REV 0413ALST  
Distributed By: Geri-Care Pharmaceuticals  
Corp. 1650 63rd Street, Brooklyn, NY 11204



3 57896 98130 3

PEEL HERE FOR MORE DRUG FACTS

## Drug Facts (continued)

**Ask a doctor or pharmacist before use if you are**

■ taking any other drug containing an NSAID (prescription or nonprescription) ■ taking a blood thinning (anticoagulant) or steroid drug ■ taking a prescription drug for diabetes, gout or arthritis

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding:

feel faint  vomit blood  have bloody or black stools  have stomach pain that does not get better  
■ an allergic reaction occurs. Seek medical help right away. ■ pain gets worse or lasts more than 10 days ■ redness or swelling is present ■ fever gets worse or lasts more than 3 days ■ any new symptoms occur ■ ringing in the ears or loss of hearing occurs

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

## Directions

■ drink a full glass of water with each dose  
■ swallow whole, do not chew or crush  
■ do not exceed recommended dose

adults and children 12 years and older

take 4-8 tablets every 4 hours, as needed, not more than 48 tablets in 24 hours, or as directed by a doctor

children under 12

ask a doctor

## Other information

■ store at room temperature 15°C - 30°C (59°F - 86°F)

**Inactive ingredients:** see carton

\*This product is not manufactured or distributed by the owner of the registered trademark Bayer®.

**STOP PEELING**