Drug Facts

Active ingredient (in each tablet) Loratadine, USP 10 mg......Antihistamine **Uses** temporarily relieves these symptoms due to hav fever or other upper respiratory allergies:

Purpose





NDC 57896-658-36

 runny nose
itchy, watery eyes sneezing
itching of the nose or throat Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness. Stop use and ask a doctor if an allergic reaction to this

product occurs. Seek medical help right away.

If pregnant or breast feeding, ask a health professional before use. Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

24-Hour Non-Drowsy*

Loratadine 10 mg Tablets

Antihistamine

Compare to active ingredient in CLARITIN®*

*When taken as directed. See Drug Facts panel

365 Tablets

Directions adults and children

12 years and over

children under 12 years of age ask a doctor consumers with liver or kidney disease ask a doctor Other information • TAMPER EVIDENT: Do not use if

imprinted seal under cap is missing or broken. • store between 20° to 25°C (68° to 77°F) • protect from light

Inactive inaredients: lactose monohydrate, magnesium stearate, microcrystalline cellulose, sodium starch glycolate

Questions or comments? call 1-800-540-3765

**This product is not manufactured or distributed by the owner of the registered trademark Claritin®.

Dist. By: GERI-CARE PHARMACEUTICALS CORP. 1295 Towbin Ave

3 57896 78836 0

1 tablet daily; not more

than 1 tablet in 24 hours

REV 0422S

Lakewood, NJ 08701