

REGULAR STRENGTH  
ENTERIC COATEDASPIRIN  
325 mg

PAIN RELIEVER (NSAID)

200 Tablets

**TAMPER EVIDENT:** Do not use if imprinted seal under cap is missing or broken.**Drug Facts****Active ingredient (in each tablet)** *Purpose*  
Aspirin 325 mg (NSAID)..... Pain reliever  
"nonsteroidal anti-inflammatory drug"**Uses** ■ for the temporary relief of minor aches and pains  
■ ask your doctor about other uses for aspirin**Warnings****Reye's Syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.**Allergy Alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling**Stomach bleeding warning:** This product contains an NSAID which may cause stomach bleeding:  
The chance is higher if you are age 60 or older■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product**Do not use** ■ if you are allergic to aspirin or any other pain reliever/fever reducer.  
■ if you ever had an allergic reaction to this product or any of its ingredients**Ask a doctor before use if** ■ the stomach bleeding warning applies to you ■ you are taking a diuretic  
■ you have a history of stomach problems, such as heartburn ■ you have:□ high blood pressure □ heart disease  
□ liver cirrhosis □ kidney disease □ asthmaDist. by: GERI-CARE PHARMACEUTICALS CORP.  
1295 Towbin Ave, Lakewood, NJ 08701

REV 1222L



3 57896192 120 0

PEEL HERE FOR MORE DRUG FACTS

Spot UV

**Drug Facts (continued)****Ask a doctor or pharmacist before use if you are** taking a prescription drug for diabetes, gout or arthritis.**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away.  
■ you experience any of the following signs of stomach bleeding: □ feel faint □ vomit blood □ have bloody or black stools  
■ have stomach pain that does not get better  
■ pain gets worse or lasts more than 10 days  
■ redness or swelling is present ■ fever gets worse or lasts more than 3 days ■ any new symptoms occur  
■ ringing in the ears or loss of hearing occurs  
These can be signs of a serious condition.**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.**Directions** ■ do not exceed recommended dose  
■ drink a full glass of water with each dose  
■ swallow whole, do not chew or crush

Take 1 to 2 tablets every 4 hours as needed, not more than 12 tablets in 24 hours, or as directed by a doctor.

Children under 12 do not use unless directed by a doctor.

**Other information** ■ store at 20-25°C (68-77°F); excursions permitted between 15-30°C (59-86°F).**Inactive ingredients:** black iron oxide, cellulose, corn starch, D&C yellow #10 lake, FD&C yellow #6 lake, hypromellose, polydextrose, polyethylene glycol, polyvinyl acetate, phthalate, propylene glycol, shellac wax, silica, simethicone, sodium alginate, sodium bicarbonate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate.**Questions or comments? 1-800-540-3765**

STOP PEELING