

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, gout or arthritis.

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: □ feel faint □ vomit blood □ have bloody or black stools ■ pain gets worse or lasts more than 10 days ■ redness or swelling is present ■ fever gets worse or lasts more than 3 days ■ any new symptoms occur ■ ringing in the ears or loss of hearing occurs

These can be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose get medical help or contact a Poison Control Center right away.

Directions ■ do not exceed recommended dose ■ drink a full glass of water with each dose ■ swallow whole, do not chew or crush

Adults and children 12 years and older: take 4-8 tablets every 4 hours, as needed, not more than 48 tablets in 24 hours, or as directed by a doctor

Children under 12: ask a doctor

Other information

■ store at 20°-25°C (68°-77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients: cellulose, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, PEG, polydextrose, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, starch, talc, titanium dioxide, triacetin, triethyl citrate, wax.

Questions or comments? 1-800-540-3765

STOP PEELING

Spot UV

GERI CARE®

NDC 57896-981-50

ADULT LOW DOSE
ENTERIC COATED

ASPIRIN 81 mg

PAIN RELIEVER (NSAID)



Compare to active ingredient
in **BAYER®** Low Dose Aspirin*

500 Tablets

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active ingredient (in each tablet) **Purpose**
Aspirin 81 mg (NSAID)*Pain reliever
*nonsteroidal anti-inflammatory drug

Uses ■ for the temporary relief of minor aches and pains
■ ask your doctor about other uses for aspirin

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ the stomach bleeding warning applies to you ■ you are taking a diuretic ■ you have a history of stomach problems, such as heartburn ■ you have: □ high blood pressure □ heart disease □ liver cirrhosis □ kidney disease □ asthma

*This product is not manufactured or distributed by the owner of the registered trademark Bayer®. Dist. by: **Geri-Care Pharmaceuticals Corp.**
1295 Towbin Avenue, Lakewood, NJ 08701

REV 0224L



3 57896 98150 1

PEEL HERE FOR MORE DRUG FACTS