Labeling Format Information:		
Font type:	Helvetica Neue Condensed	
Barline	2.0 pt	
Hairline	0.5 pt	
Drug facts:	10.5 pt	
Drug facts (continued):	9.5 pt	
Header:	8.5 pt	
Subheader:	6.5 pt	
Body text:	6.5 pt	
Bullets:	6.5 pt	

Drug Facts

Active ingredient — **Purpose**

Diclofenac sodium USP (NSAID*) 1% (equivalent to 0.93% diclofenac) ..... Arthritis pain reliever

*nonsteroidal anti-inflammatory drug

Uses

- for the temporary relief of arthritis pain ONLY in the following areas:
  - hand, wrist, elbow (upper body areas)
  - foot, ankle, knee (lower body areas)
- this product may take up to 7 days to work for arthritis pain; it is not for immediate relief. If no pain relief in 7 days, stop use.

Warnings

For external use only

Allergy alert: Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- blisters
- facial swelling
- shock
- rash

if an allergic reaction occurs, stop use and seek medical help right away.

↓ Lift Flap for Important Directions & Warnings ↓

LIFT HERE ↓

Drug Facts (continued)

Liver warning: This product contains diclofenac. Liver damage may occur if you apply

- more or for a longer time than directed
- when using other drugs containing diclofenac

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is small but higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- apply more or for longer than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal.

The risk is higher if you use more than directed or for longer than directed.

Use enclosed dosing card to measure a dose

Diclofenac Sodium Topical Gel, 1%
Arthritis Pain
NSAID - Arthritis Pain Reliever

Upper body dose 2.25 inches long (2 grams)

Lower body dose 4.5 inches long (4 grams)

Dosing Card (NOT ACTUAL SIZE)
Read product User Guide before use
Not for immediate pain relief

How to Use

- ☑ For arthritis pain only
- ☑ Use 4 times a day
- ☑ May take up to 7 days to work for your arthritis pain
- ☑ For use on no more than 2 body areas
- ☑ Use up to 21 days unless directed by your doctor

Save carton, dosing card, and User Guide for full product uses, directions and warnings.

TAMPER EVIDENT STATEMENT: DO NOT USE IF SAFETY SEAL ON TUBE IS BROKEN OR MISSING

ORIGINAL PRESCRIPTION STRENGTH

GERI CARE®

NDC 57896-147-01

Compare to the active ingredient in Voltaren® Arthritis Pain**

Diclofenac Sodium Topical Gel, 1%

Arthritis Pain

NSAID - Arthritis Pain Reliever

NET WT. 3.53 oz (100 g)

- For Daily Treatment of Arthritis Pain
- Anti-Inflammatory

For external use only

Lot:
Exp:

UVZ

P1052541 LM-5353

Medicated Gel Clinically Proven to Relieve Arthritis Pain

Distributed by:
GERI-CARE PHARMACEUTICALS CORP.
1295 Towbin Avenue
Lakewood, NJ 08701

Made in India
Code: TS/DRUGS/58039/2020

Rev. 07/23

TREATMENT AREAS

Not for use on any other body area (such as back, hip, or shoulder)

**This product is not manufactured or distributed by Novartis Corporation, owner of the registered trademark Voltaren®.

Drug Facts (continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Use up to 21 days unless directed by your doctor

Not for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.

Daily	Per Dose
For your arthritis pain: • Use 4 times per day every day • Do not use on more than 2 body areas at the same time	**Use ENCLOSED DOSING CARD to measure a dose** • For each upper body area (hand, wrist, or elbow) - Squeeze out 2.25 inches (2 grams) • For each lower body area (foot, ankle or knee) - Squeeze out 4.5 inches (4 grams)

Read the enclosed User Guide for complete instructions:

- use only as directed

Drug Facts (continued)

- do not use more than directed or for longer than directed
- apply only to clean, dry skin that does not have any cuts, open wounds, infections or rashes
- do not apply in same area as any other product
- do not apply with external heat such as heating pad
- do not apply a bandage over the treated area
- store ENCLOSED DOSING CARD with your diclofenac sodium topical gel arthritis pain. The dosing card is re-usable.

Other information

- store at 20° to 25°C (68° to 77°F). Keep from freezing.
- read all product information before using. Keep the dosing card, this carton and accompanying User Guide for important information.

Inactive ingredients

carbomer homopolymer Type C, cocoyl caprylocaprate, fragrance, isopropyl alcohol, mineral oil, polyoxyl 20 cetostearyl ether, propylene glycol, purified water, strong ammonia solution

Questions or comments? call 1-855-274-4122

3 57896 14701 3