

**TAMPER-EVIDENT: DO NOT USE IF
IMPRINTED NECKBAND IS MISSING
OR BROKEN.**

Drug Facts

Active ingredients (in each 10 mL dose)

Aluminum hydroxide (equivalent to dried gel, USP) 800 mg.....	Antacid
Magnesium hydroxide 800 mg.....	Antacid
Simethicone 80 mg	Antigas

Purposes

Uses relieves • heartburn • sour stomach
• acid indigestion • the symptoms referred to as gas

Warnings

Ask a doctor before use if you have

- kidney disease • a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions • shake well before use

- adults and children 12 years of age and older: take 10 mL to 20 mL (1 to 2 doses) between meals as needed, at bedtime, or as directed by a doctor
- do not take more than 60 mL (6 doses) in a 24 hour period or use the maximum dosage for more than 2 weeks
- children under 12 years of age: ask a doctor
- measure with dosing cup provided
- mL = milliliter

Other information

- **each 10 mL dose contains:** magnesium 340 mg, sodium 10 mg
- do not freeze • store at room temperature tightly closed

Inactive ingredients benzyl alcohol, carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose, purified water, saccharin sodium, sorbitol solution

Questions or comments? 1-800-540-3765

*This product is not manufactured or distributed by the owner of the registered trademark Mylanta® Maximum Strength.

910-06142-0 REV 0524



Distributed by
Geri-Care Pharmaceuticals
1295 Towbin Ave.
Lakewood, NJ 08701