

Drug Facts

Active ingredients (in each 5 mL teaspoonful)

Aluminum hydroxide 254 mg	Antacid
Magnesium carbonate 237.5 mg	Antacid

Uses relieves • acid indigestion • heartburn • sour stomach
• upset stomach associated with these symptoms

Warnings

Ask a doctor before use if you have • kidney disease
• a magnesium-restricted diet • a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescription drugs.

When using this product • laxative effect may occur
• do not use maximum dosage for more than 2 weeks
• do not take more than 16 teaspoonfuls in 24 hours

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222.

Directions

- shake well before use
- adults and children 12 years and older take 2-4 teaspoonsful four times a day or as directed by a doctor
- children under 12 years: consult a doctor
- take after meals and at bedtime
- dispense product only by spoon or other measuring device

Other information

- **each 5 mL teaspoonful contains:** magnesium 70 mg, sodium 16 mg
- store at room temperature • protect from freezing • keep tightly closed
- **TAMPER-EVIDENT:** Do not use this product if inner foil seal over mouth of the bottle is broken or missing.

Inactive ingredients benzyl alcohol, edetate disodium, flavor, glycerin, purified water, saccharin sodium, simethicone emulsion, sodium alginate, sorbitol solution, xanthan gum

Questions or comments? 1-800-540-3765

*This product is not manufactured or distributed by GlaxoSmithKline, owner of the registered trademark Gaviscon® Extra Strength.

Distributed by
Geri-Care Pharmaceuticals
1295 Towbin Ave.
Lakewood, NJ 08701

910-17222-0
REV 0624

