

Drug Facts

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
NECKBAND IS MISSING OR BROKEN.**

Active ingredient (in each 15 mL tablespoonful) Purpose

Magnesium hydroxide 1200 mg Saline laxative

Use

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are taking a prescription drug.
This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product.
These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or
contact a Poison Control Center (1-800-222-1222) right away.

Directions

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses,
or as directed by a doctor
- follow each dosage with a full glass (8 oz) of liquid

age	dose
adults and children 12 years and over	2 to 4 tablespoonfuls
children 6 to 11 years	1 to 2 tablespoonfuls
children under 6 years	ask a doctor

Other information

- each 15 mL contains: magnesium 500 mg, sodium 6 mg
- do not freeze • store at room temperature tightly closed

Inactive ingredients carboxymethylcellulose sodium, citric acid,
D&C red no. 28, flavor, glycerin, microcrystalline cellulose, purified water,
sodium citrate, sodium hypochlorite, sucrose, xanthan gum

Questions or comments? 1-800-540-3765

Dist. By: **Geri-Care Pharmaceuticals Corp.**
1295 Towbin Avenue, Lakewood, NJ 08701
1-800-540-3765

REV 0424
910-19512-0

