

Drug Facts (continued)

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions: Chew tablets thoroughly before swallowing.

This product does not contain directions or complete warnings for adults.

Follow dosage on table or use as doctor directs. If possible use weight to dose; otherwise use age. Give every 4 hours as needed; do not give more than 5 doses in 24 hours. Administer only under adult supervision.

DO NOT EXCEED DOSE SHOWN.

See Overdose Warning.

Weight (lbs)	Under 24	24-35	36-47	48-59	60-71	72-95
Age	Under 2	2-3	4-5	6-8	9-10	11
Tablets	Ask a Doctor	2	3	4	5	6

Other information

• **TAMPER EVIDENT: Do not use if imprinted seal under cap is missing or broken.**

- store at 20-25°C (68-77°F); avoid high humidity
- Keep tightly closed
- Phenylketonurics: contains phenylalanine 5mg per tablet.

Inactive ingredients:

Aspartame (contains phenylalanine), cellulose, D&C Red #27, Dextrose, FD&C Yellow #6, fruit flavor, magnesium stearate, Prosweet, stearic acid, sugar.

Questions or comments? 1-800-540-3765

STOP PEELING

HEALTH STAR[®]

NDC 57896-148-03

CHILDREN'S CHEWABLE ACETAMINOPHEN 80 mg

PAIN RELIEVER/ FEVER REDUCER

FRUIT FLAVOR
CHEWABLE TABLETS

30 Tablets 80 mg each

TAMPER EVIDENT: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 80 mg.....Pain Reliever / Fever Reducer

Uses • temporarily relieves minor aches and pains
• temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount

• with other drugs containing acetaminophen
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

DIST. BY: GRI-CARE PHARMACEUTICALS CORP.
1295 Towbin Ave, Lakewood, NJ 08701

REV 0224Ad
3 57896 14803 4

PEEL HERE FOR MORE DRUG FACTS