

NDC 57896-317-03



ORIGINAL STRENGTH

FAMOTIDINE TABLETS, USP 10 mg

ACID REDUCER

Just One Tablet
Prevents & Relieves Heartburn
Due To Acid Indigestion

ACTUAL SIZE



Compare to active ingredient in
Original Strength **PEPCID AC**®*

30
TABLETS

Do not use if printed foil seal on bottle is broken or missing.

Drug Facts

Active ingredient **Purpose**
(in each tablet)

Famotidine, USP 10 mg.....Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

■ with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

Drug Facts (continued)

- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days

if pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20°C - 25°C (68°F - 77°F) ■ protect from moisture

NDC 57896-317-03



ORIGINAL STRENGTH

FAMOTIDINE TABLETS USP 10 mg, ACID REDUCER

Distributed by: **Gerl-Care Pharmaceuticals Corp.**
1295 Towbin Avenue, Lakewood, NJ 08701
Product of India M.L. No.: 02/SKL/AP/2015/F/R

*This product is not manufactured or distributed by the owner of the registered trademark **PEPCID AC**®, nor by the distributor of **PEPCID AC**®. 800228

Drug Facts (continued)

Inactive ingredients

Microcrystalline Cellulose, Pre-gelatinized Starch, Sodium Starch Glycolate, Hydroxypropyl Cellulose, Magnesium Stearate, Hypromellose, Titanium Dioxide, Triacetin, Talc, Macrogol

Questions or comments?

1-800-540-3765 (toll-free)

JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages

TIPS FOR MANAGING HEARTBURN

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

REV 03/24

